=> file medline caplus biosis biotechds scisearch embase

COST IN U.S. DOLLARS SINCE FILE TOTAL

ENTRY SESSION 0.42 0.42

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 11:44:23 ON 14 JAN 2003

FILE 'CAPLUS' ENTERED AT 11:44:23 ON 14 JAN 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 11:44:23 ON 14 JAN 2003 COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC. (R)

FILE 'BIOTECHDS' ENTERED AT 11:44:23 ON 14 JAN 2003

COPYRIGHT (C) 2003 THOMSON DERWENT AND INSTITUTE FOR SCIENTIFIC INFORMATION

FILE 'SCISEARCH' ENTERED AT 11:44:23 ON 14 JAN 2003

COPYRIGHT (C) 2003 Institute for Scientific Information (ISI) (R)

FILE 'EMBASE' ENTERED AT 11:44:23 ON 14 JAN 2003

COPYRIGHT (C) 2003 Elsevier Science B.V. All rights reserved.

=> s monooxygenase and oxidation

L1 7515 MONOOXYGENASE AND OXIDATION

=> dup rem 11

PROCESSING IS APPROXIMATELY 19% COMPLETE FOR L1

PROCESSING IS APPROXIMATELY 52% COMPLETE FOR L1

PROCESSING IS APPROXIMATELY 72% COMPLETE FOR L1

PROCESSING IS APPROXIMATELY 95% COMPLETE FOR L1

PROCESSING COMPLETED FOR L1

L2 4386 DUP REM L1 (3129 DUPLICATES REMOVED)

=> s 12 and cytochrome P-450cam

4 FILES SEARCHED...

L3 28 L2 AND CYTOCHROME P-450CAM

=> s 13 and hologenated aromatic

L4 0 L3 AND HOLOGENATED AROMATIC

=> s 13 and halogenated aromatic

L5 0 L3 AND HALOGENATED AROMATIC

=> s 13 and halo aromatic

L6 0 L3 AND HALO AROMATIC

=> s 13 and aromatic

L7 5 L3 AND AROMATIC

=> d 17 1-5 ibib ab

L7 ANSWER 1 OF 5 MEDLINE

ACCESSION NUMBER: 97163846 MEDLINE

DOCUMENT NUMBER: 97163846 PubMed ID: 9010601

TITLE: A structure-based model for cytochrome P450cam-

putidaredoxin interactions.

AUTHOR: Pochapsky T C; Lyons T A; Kazanis S; Arakaki T; Ratnaswamy

G

CORPORATE SOURCE: Department of Chemistry, Brandeis University, Waltham, MA

02254-9110, USA.

CONTRACT NUMBER: RO1-GM-44191 (NIGMS)

BIOCHIMIE, (1996) 78 (8-9) 723-33. SOURCE:

Journal code: 1264604. ISSN: 0300-9084.

PUB. COUNTRY:

France

DOCUMENT TYPE:

Journal; Article; (JOURNAL ARTICLE)

LANGUAGE:

English

FILE SEGMENT:

Priority Journals

ENTRY MONTH:

199707

ENTRY DATE:

Entered STN: 19970721

Last Updated on STN: 19970721 Entered Medline: 19970708

Putidaredoxin (Pdx) is a Fe2S2 ferredoxin which acts as the physiological AB reductant of cytochrome P-450cam (CYP101). A model for the solution structure of oxidized Pdx has been determined using NMR methods (Pochapsky et al (1994) Biochemistry 33, 6424-6432). 1H-15N correlations and redox-dependent amide exchange rates have also been described (Lyons et al (1996) Protein Sci 5, 627-639). Data obtained from mutagenesis and kinetic measurements concerning the interactions of Pdx and CYP101 are summarized. A model for the structure of the homologous ferredoxin adrenodoxin (Adx) is also described, and data concerning Adx activity are discussed in relation to this structure. The structures of Pdx and CYP101 were used as starting points for molecular modeling and molecular dynamics simulations. Close approach between the metal centers of the two proteins and interaction between aromatic residues on the surfaces of the proteins are premised. The resulting complex exhibits three intermolecular salt bridges, five intermolecular hydrogen bonds and a 12 A distance between the metal centers. The first direct observations of interaction between Pdx and CYP101 (by two-dimensional NMR of 15N-labeled Pdx in solution with CYP101) are described. The results of the NMR experiments indicate that conformational gating of the electron transfer complex between CYP101 and Pdx may be important.

ANSWER 2 OF 5 CAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER:

2002:220824 CAPLUS

DOCUMENT NUMBER:

136:259213

TITLE:

Screening method for oxygenase enzymes using aromatic substrates which are converted to spectrochemically detected polymeric oxygenated

compounds by a coupling enzyme Arnold, Frances H.; Joo, Hyun

PATENT ASSIGNEE(S):

California Institute of Technology, USA

SOURCE:

PCT Int. Appl., 125 pp.

CODEN: PIXXD2

DOCUMENT TYPE:

INVENTOR(S):

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT:

PATENT INFORMATION:

```
APPLICATION NO. DATE
                                   KIND
                                              DATE
        PATENT NO.
                                                                   . _____
                                    _---
                                                                 WO 2000-US28768 20001013
       WO 2002022861
                                   A1
                                              20020321
              W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
              CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DY, ES, ET, EB, CB, CB, CB, TE, LT, LU, MC, NI, PT, SE, BF, BJ
                     DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
                     CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
                                              20020326
                                                                      AU 2000-80285
                                                                                                    20001013
       AU 2000080285
                                     Α5
PRIORITY APPLN. INFO.:
                                                                  US 2000-661093 A1 20000913
                                                                  WO 2000-US28768 W 20001013
```

A method is provided for detecting the presence of an oxygenated compd. AΒ

which is produced when a substrate is reacted with an oxygenase for the substrate. The method involves reacting a coupling enzyme with the oxygenated compd. to form a polymeric oxygenated compd. which is fluorescent or luminescent. Measurement of the fluorescence or luminescence of the polymeric oxygenated compd. provides indirect detection of the oxygenated compd. produced by reaction of the oxygenase with the substrate. The method is carried out in a whole cell environment wherein the cell is transformed to express both the oxygenase being screened and the coupling enzyme. The method can be used to measure the activity of monooxygenases and dioxygenases on arom. substrates. Thus, for example, the activity of cytochrome P

450cam in Escherichia coli is checked by measuring the conversion of naphthalene to a hydroxylated product (e.g., 1-naphthol, 2-naphthol) which emits a blue fluorescence when exogenously added horseradish peroxidase polymerizes the product. The method is amenable to large scale screening of enzyme mutants to isolate those with max. oxygenase activity. Thus, a screening strategy with high throughput fluorescence image anal. was implemented in order to identify bacterial clones expressing improved hydroxylating enzymes. Mutants of P 450cam with improved activity on naphthalene (or 3-phenylpropionate) and H2O2 are identified.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER:

2000:911441 CAPLUS

DOCUMENT NUMBER:

134:68048

TITLE:

Analogs of a cytochrome P450 of Pseudomonas putida

with improved catalytic action aromatic

halohydrocarbons for use in bioremediation of soil

INVENTOR(S):

Wong, Luet Lok; Jones, Jonathan Peter

PATENT ASSIGNEE(S):

Isis Innovation Limited, UK

SOURCE:

PCT Int. Appl., 42 pp. CODEN: PIXXD2

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

```
APPLICATION NO. DATE
     PATENT NO.
                     KIND DATE
                                            WO 2000-GB2379 20000619
     WO 2000078973
                      A1 20001228
         W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
             CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
             LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,
             SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
             YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
         RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
             DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
             CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
                                            EP 2000-942200 20000619
                       A1
                            20020327
     EP 1190067
         R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
             IE, SI, LT, LV, FI, RO
                                          GB 1999-14373
PRIORITY APPLN. INFO.:
                                                            A 19990618
                                          WO 2000-GB2379
                                                            W 20000619
```

AB Analaogs of cytochrome P 450cam of

Pseudomonas putida that have improved catalytic activity against heavily halogenated arom. hydrocarbons and that may be of use in the reclamation of soils contaminated with polychlorinated biphenyls. In particular, alterations in the substrate pocket that increase the vol. available for bulky polyhalogenated aroms. are described. Prepn. of a series of analogs of the gene camC cytochrome P 450 with increased activity towards polychlorinated biphenyls is demonstrated. A fusion protein of

putidaredoxin and putidaredoxin reductase that can be used as a cofactor is also described.

REFERENCE COUNT:

THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 5 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: DOCUMENT NUMBER:

1994:133100 BIOSIS PREV199497146100

TITLE:

Putative functions of phenylalanine-350 of Pseudomonas

putida cytochrome P-450-cam.

AUTHOR(S):

Yasukochi, Takanori; Okada, Osamu; Hara, Takayuki; Sagara,

Yasuhiro (1); Sekimizu, Kazuhisa; Horiuchi, Tadao

CORPORATE SOURCE:

(1) Dep. Med. Biol., Kochi Med. Sch., Okoh-cho, Nankoku,

Kochi 783 Japan

SOURCE:

Biochimica et Biophysica Acta, (1994) Vol. 1204, No. 1, pp.

84-90.

ISSN: 0006-3002.

DOCUMENT TYPE:

Article English

LANGUAGE: Cytochrome P-450-cam hydroxylates d-camphor, using molecular oxygen and reducing equivalents transferred via putidaredoxin. We constructed mutant genes in which Phe-350 of P-450-cam was replaced by Leu, Tyr, or His by site-directed mutagenesis, expressed them in Escherichia coli, purified the mutant proteins, and compared their enzymic properties with those of the wild type P-450-cam. NADH oxidation rate of the Tyr mutant in the reconstituted system with putidaredoxin and putidaredoxin reductase was similar to that of the wild type enzyme, while the Leu mutant and the His mutant showed 67% and 17% activity of that of the wild type, respectively. The affinities of these mutant proteins for camphor and the oxidized form of putidaredoxin were much the same as those of the wild type protein. Rate constants for the reduction reaction of P-450-cam by reduced putidaredoxin, a physiological electron donor for P-450-cam, of Tyr and His mutants were much the same as that of the wild type enzyme, whereas the Leu mutant showed approx. half that of the wild type. Thus, the aromatic ring of Phe-350 of P-450-cam probably contributes to enhancing efficiency of the electron transfer yet does not seem to be essential for the reaction.

L7 ANSWER 5 OF 5 SCISEARCH COPYRIGHT 2003 ISI (R)

ACCESSION NUMBER: 97:87565 SCISEARCH

THE GENUINE ARTICLE: WD134

TITLE:

Gene organization and low regiospecificity in aromatic-ring hydroxylation of a benzene

monooxygenase of Pseudomonas aeruginosa JI104

AUTHOR: CORPORATE SOURCE:

Kitayama A (Reprint); Suzuki E; Kawakami Y; Nagamune T UNIV TOKYO, GRAD SCH ENGN, DEPT CHEM & BIOTECHNOL, BUNKYO KU, 7-3-1 HONGO, TOKYO 113, JAPAN (Reprint); INST RES &

INNOVAT, DEPT BIOTECHNOL, KASHIWA, CHIBA 277, JAPAN

COUNTRY OF AUTHOR: JAPAN

SOURCE:

JOURNAL OF FERMENTATION AND BIOENGINEERING, (20 JAN 1996)

Vol. 82, No. 5, pp. 421-425.

Publisher: SOC FERMENTATION BIOENGINEERING, JAPAN, OSAKA UNIV, FACULTY ENGINEERING, 2-1 YAMADAOKA, SUITA, OSAKA

565, JAPAN. ISSN: 0922-338X.

DOCUMENT TYPE:

REFERENCE COUNT:

Article; Journal

FILE SEGMENT: LIFE; AGRI LANGUAGE: English

ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

AB A novel benzene monooxygenase gene cluster has been cloned from Pseudomonas aeruginosa J104 which was isolated from soil as a benzene degrader. The nucleotide sequence of this gene cluster was found to be

highly homologous to those of other toluene monocxygenase gene clusters. This multicomponent monocxygenase also has the capability to catalyze the hydroxylation of various alkylated aromatic hydrocarbons. The low regiospecific hydroxylation was observed when toluene, o-xylene, ethyl benzene and n-propyl benzene were used as substrates.

```
=> s 13 and halogenated substrate
             O L3 AND HALOGENATED SUBSTRATE
=> d his
     (FILE 'HOME' ENTERED AT 11:43:23 ON 14 JAN 2003)
     FILE 'MEDLINE, CAPLUS, BIOSIS, BIOTECHDS, SCISEARCH, EMBASE' ENTERED AT
     11:44:23 ON 14 JAN 2003
           7515 S MONOOXYGENASE AND OXIDATION
L1
L2
           4386 DUP REM L1 (3129 DUPLICATES REMOVED)
L3
             28 S L2 AND CYTOCHROME P-450CAM
              0 S L3 AND HOLOGENATED AROMATIC
L4
L5
              0 S L3 AND HALOGENATED AROMATIC
              0 S L3 AND HALO AROMATIC
L6
              5 S L3 AND AROMATIC
L7
              0 S L3 AND HALOGENATED SUBSTRATE
T.8
=> s 12 and halogenated aromatic substrate
   2 FILES SEARCHED...
             O L2 AND HALOGENATED AROMATIC SUBSTRATE
=> s 13 and dichlorobenzene
             1 L3 AND DICHLOROBENZENE
=> d 110 ibib ab
L10 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2003 ACS
                          2000:74430 CAPLUS
ACCESSION NUMBER:
DOCUMENT NUMBER:
                          132:141329
                          The oxidation of polychlorinated benzenes by
TITLE:
                          genetically engineered cytochrome P450cam: potential
                          applications in bioremediation
                         Jones, Jonathan P.; O'Hare, Ellen J.; Wong, Luet-Lok
Dep. Chem., Inorganic Chem. Lab., University of
AUTHOR(S):
CORPORATE SOURCE:
                          Oxford, Oxford, OX1 3QR, UK
                          Chemical Communications (Cambridge) (2000), (3),
SOURCE:
                          247-248
                          CODEN: CHCOFS; ISSN: 1359-7345
                          Royal Society of Chemistry
PUBLISHER:
DOCUMENT TYPE:
                          Journal
LANGUAGE:
                          English
     Polychlorinated arom. compds. are persistent environmental pollutants.; we
     describe here. Redesign and engineering of the heme monooxygenase
     , cytochrome P 450cam, to oxidize these
     compds. efficiently to chlorinated phenols which are readily degraded by
     many microorganisms, thus provides a basis for novel bioremediation
     systems for these inert compds., are described.
                               THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS
REFERENCE COUNT:
                         13
                                RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT
```

=> s 13 and (benzene or biphenyl) L11 5 L3 AND (BENZENE OR BIPHENYL) L11 ANSWER 1 OF 5

MEDLINE

ACCESSION NUMBER:

2002612268 MEDLINE

DOCUMENT NUMBER:

22241836 PubMed ID: 12114516

TITLE:

Crystal structure of the F87W/Y96F/V247L mutant of

cvtochrome P-450cam with

1,3,5-trichlorobenzene bound and further protein

engineering for the oxidation of

pentachlorobenzene and hexachlorobenzene.

AUTHOR:

Chen Xuehui; Christopher Alexandra; Jones Jonathan P; Bell

Stephen G; Guo Qing; Xu Feng; Rao Zihe; Wong Luet-Lok

CORPORATE SOURCE:

Laboratory of Structural Biology, Department of Biological Science and Technology & Ministry of Education Laboratory of Protein Science, Tsinghua University, Beijing 100084,

SOURCE:

JOURNAL OF BIOLOGICAL CHEMISTRY, (2002 Oct 4) 277 (40)

37519-26.

Journal code: 2985121R. ISSN: 0021-9258.

PUB. COUNTRY:

United States

DOCUMENT TYPE:

Journal; Article; (JOURNAL ARTICLE)

LANGUAGE:

English

FILE SEGMENT:

Priority Journals

OTHER SOURCE:

PDB-1J51 200211

ENTRY MONTH: ENTRY DATE:

Entered STN: 20021010

Last Updated on STN: 20030105

Entered Medline: 20021120

We reported previously that the F87W/Y96F/V247L mutant of AB

cytochrome P-450cam (CYP101) from Pseudomonas putida catalyzed the rapid oxidation of lightly chlorinated

benzenes, but pentachlorobenzene oxidation was slow (Jones, J. P., O'Hare, E. J., and Wong, L. L. (2001) Eur. J. Biochem. 268, 1460-1467). In the present work, we determined the crystal structure of this mutant with bound 1,3,5-trichlorobenzene. The substrate was bound to crystallographically independent CYP101 molecules in at least three different orientations, which were distinguished by the angle between the benzene ring and the porphyrin, and one orientation contained an Fe-Cl interaction. In another orientation, the substrate was almost parallel to the heme, with a C-H bond closest to the iron. The enzyme/substrate contacts suggested that the L244A mutation should promote the binding of pentachlorobenzene and hexachlorobenzene by creating space to accommodate the extra chlorines. The F87W/Y96F/L244A/V247L mutant thus designed was found to oxidize pentachlorobenzene at a rate of 82.5 nmol

200-fold, to 2.0 min(-1). Both substrates are oxidized to pentachlorophenol, which is degraded by micro-organisms. In principle, the F87W/Y96F/L244A/V247L mutant could have applications in the bioremediation of polychlorinated benzenes.

 $(nmol\ CYP101)\ (-1)\ min\ (-1)$, 45 times faster than the F87W/Y96F/V247L parent

L11 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER:

2002:220824 CAPLUS

mutant. The rate of hexachlorobenzene oxidation was increased

DOCUMENT NUMBER:

136:259213

TITLE:

Screening method for oxygenase enzymes using aromatic substrates which are converted to spectrochemically detected polymeric oxygenated compounds by a coupling

enzyme

INVENTOR(S):

Arnold, Frances H.; Joo, Hyun

PATENT ASSIGNEE(S):

California Institute of Technology, USA

SOURCE:

PCT Int. Appl., 125 pp. CODEN: PIXXD2

DOCUMENT TYPE:

Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

```
APPLICATION NO. DATE
    PATENT NO.
                   KIND DATE
                                         -----
    _____ ___
                          <del>------</del>
                                        WO 2000-US28768 20001013
    WO 2002022861
                    A1 20020321
        W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
            CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL,
            IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
            MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG,
            SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW,
            AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
        RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
            DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
            CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
                                        AU 2000-80285
                                                          20001013
    AU 2000080285
                     A5
                          20020326
                                      US 2000-661093 A1 20000913
PRIORITY APPLN. INFO .:
                                      WO 2000-US28768 W 20001013
```

Amethod is provided for detecting the presence of an oxygenated compd. which is produced when a substrate is reacted with an oxygenase for the substrate. The method involves reacting a coupling enzyme with the oxygenated compd. to form a polymeric oxygenated compd. which is fluorescent or luminescent. Measurement of the fluorescence or luminescence of the polymeric oxygenated compd. provides indirect detection of the oxygenated compd. produced by reaction of the oxygenase with the substrate. The method is carried out in a whole cell environment wherein the cell is transformed to express both the oxygenase being screened and the coupling enzyme. The method can be used to measure the activity of monooxygenases and dioxygenases on arom. substrates. Thus, for example, the activity of cytochrome P

450cam in Escherichia coli is checked by measuring the conversion of naphthalene to a hydroxylated product (e.g., 1-naphthol, 2-naphthol) which emits a blue fluorescence when exogenously added horseradish peroxidase polymerizes the product. The method is amenable to large scale screening of enzyme mutants to isolate those with max. oxygenase activity. Thus, a screening strategy with high throughput fluorescence image anal. was implemented in order to identify bacterial clones expressing improved hydroxylating enzymes. Mutants of P 450cam with improved activity on naphthalene (or 3-phenylpropionate) and H2O2 are identified.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2003 ACS ACCESSION NUMBER: 2000:911441 CAPLUS

DOCUMENT NUMBER: 134:68048

TITLE: Analogs of a cytochrome P450 of Pseudomonas putida

with improved catalytic action aromatic

halohydrocarbons for use in bioremediation of soil

INVENTOR(S): Wong, Luet Lok; Jones, Jonathan Peter

PATENT ASSIGNEE(S): Isis Innovation Limited, UK

SOURCE: PCT Int. Appl., 42 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.			KI	ND	DATE			A.	PPLI	CATI	ои ис	ο.	DATE			
WO 2000078973		A	1	2000	001228 WO 2000-GB2379		9	20000619								
w:	ΑE,	AG,	AL,	AM,	AT,	ΑU,	ΑZ,	BA,	BB,	BG,	BR,	BY,	ΒZ,	CA,	CH,	CN,
	CR,	CU,	CZ,	DE,	DK,	DM,	DZ,	EE,	ES,	FI,	GB,	GD,	GE,	GH,	GM,	HR,

```
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,
             SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
         RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
              CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
                                             EP 2000-942200 20000619
                             20020327
                        A1
            AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
             IE, SI, LT, LV, FI, RO
PRIORITY APPLN. INFO.:
                                           GB 1999-14373
                                                             A 19990618
                                           WO 2000-GB2379
                                                             W 20000619
AΒ
     Analaogs of cytochrome P 450cam of
     Pseudomonas putida that have improved catalytic activity against heavily
     halogenated arom. hydrocarbons and that may be of use in the reclamation
     of soils contaminated with polychlorinated biphenyls. In particular,
     alterations in the substrate pocket that increase the vol. available for
     bulky polyhalogenated aroms. are described. Prepn. of a series of analogs
     of the gene camC cytochrome P 450 with increased activity towards
     polychlorinated biphenyls is demonstrated. A fusion protein of
     putidaredoxin and putidaredoxin reductase that can be used as a cofactor
     is also described.
REFERENCE COUNT:
                          7
                                 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
                                 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT
L11 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2003 ACS
                          2000:74430 CAPLUS
ACCESSION NUMBER:
DOCUMENT NUMBER:
                          132:141329
TITLE:
                          The oxidation of polychlorinated benzenes by
                          genetically engineered cytochrome P450cam: potential
                          applications in bioremediation
                          Jones, Jonathan P.; O'Hare, Ellen J.; Wong, Luet-Lok
AUTHOR(S):
                          Dep. Chem., Inorganic Chem. Lab., University of
CORPORATE SOURCE:
                          Oxford, Oxford, OX1 3QR, UK
                          Chemical Communications (Cambridge) (2000), (3),
SOURCE:
                          247-248
                          CODEN: CHCOFS; ISSN: 1359-7345
PUBLISHER:
                          Royal Society of Chemistry
DOCUMENT TYPE:
                          Journal
                          English
LANGUAGE:
     Polychlorinated arom. compds. are persistent environmental pollutants.; we
     describe here. Redesign and engineering of the heme monooxygenase
     , cytochrome P 450cam, to oxidize these
     compds. efficiently to chlorinated phenols which are readily degraded by
     many microorganisms, thus provides a basis for novel bioremediation
     systems for these inert compds., are described.
REFERENCE COUNT:
                          13
                                 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS
                                 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT
L11 ANSWER 5 OF 5 SCISEARCH
                                COPYRIGHT 2003 ISI (R)
                      97:87565
                                SCISEARCH
ACCESSION NUMBER:
THE GENUINE ARTICLE: WD134
TITLE:
                      Gene organization and low regiospecificity in
                      aromatic-ring hydroxylation of a benzene
                      monooxygenase of Pseudomonas aeruginosa JI104
                      Kitayama A (Reprint); Suzuki E; Kawakami Y; Nagamune T
AUTHOR:
                      UNIV TOKYO, GRAD SCH ENGN, DEPT CHEM & BIOTECHNOL, BUNKYO
CORPORATE SOURCE:
                      KU, 7-3-1 HONGO, TOKYO 113, JAPAN (Reprint); INST RES &
                      INNOVAT, DEPT BIOTECHNOL, KASHIWA, CHIBA 277, JAPAN
COUNTRY OF AUTHOR:
                      JAPAN
                      JOURNAL OF FERMENTATION AND BIOENGINEERING, (20 JAN 1996)
SOURCE:
                      Vol. 82, No. 5, pp. 421-425.
```

Publisher: SOC FERMENTATION BIOENGINEERING, JAPAN, OSAKA

UNIV, FACULTY ENGINEERING, 2-1 YAMADAOKA, SUITA, OSAKA

565, JAPAN.

ISSN: 0922-338X. Article; Journal

DOCUMENT TYPE: FILE SEGMENT:

LIFE; AGRI English

LANGUAGE:

REFERENCE COUNT: 20

ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

A novel benzene monooxygenase gene cluster has been

cloned from Pseudomonas aeruginosa J104 which was isolated from soil as a benzene degrader. The nucleotide sequence of this gene cluster was

found to be highly homologous to those of other toluene

monooxygenase gene clusters. This multicomponent

monooxygenase also has the capability to catalyze the

hydroxylation of various alkylated aromatic hydrocarbons. The low regiospecific hydroxylation was observed when toluene, o-xylene, ethyl

benzene and n-propyl benzene were used as substrates.

=> s 13 and (chlorine)

L12 1 L3 AND (CHLORINE)

=> d 112

L12 ANSWER 1 OF 1 MEDLINE

93333178 MEDLINE ΑN

93333178 PubMed ID: 7763853 DN

Cosubstrate effects in reductive dehalogenation by Pseudomonas putida G786 TI expressing cytochrome P-450CAM.

ΑU Logan M S; Newman L M; Schanke C A; Wackett L P

Gray Freshwater Biological Institute, University of Minnesota, Navarre CS 55392.

NC GM41235 (NIGMS)

BIODEGRADATION, (1993) 4 (1) 39-50. SO Journal code: 9100834. ISSN: 0923-9820.

CY Netherlands

DΤ Journal; Article; (JOURNAL ARTICLE)

LΑ English

Biotechnology FS

EM 199309

ED Entered STN: 19950809

> Last Updated on STN: 19970203 Entered Medline: 19930901

=> d 112 ibib ab

L12 ANSWER 1 OF 1 MEDLINE

MEDLINE ACCESSION NUMBER: 93333178

DOCUMENT NUMBER: 93333178 PubMed ID: 7763853

TITLE: Cosubstrate effects in reductive dehalogenation by

Pseudomonas putida G786 expressing cytochrome

Logan M S; Newman L M; Schanke C A; Wackett L P AUTHOR:

CORPORATE SOURCE: Gray Freshwater Biological Institute, University of

Minnesota, Navarre 55392.

CONTRACT NUMBER: GM41235 (NIGMS)

SOURCE: BIODEGRADATION, (1993) 4 (1) 39-50.

Journal code: 9100834. ISSN: 0923-9820.

PUB. COUNTRY: Netherlands

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Biotechnology ENTRY MONTH:

199309

ENTRY DATE:

Entered STN: 19950809

Last Updated on STN: 19970203 Entered Medline: 19930901

AB Cytochrome P-450CAM was shown to be the

primary catalyst mediating reductive dehalogenation of polychlorinated ethanes by Pseudomonas putida G786. Under anaerobic conditions, the enzyme catalyzed reductive elimination reactions in vivo with the substrates hexachloroethane, pentachloroethane, and 1,1,1,2-tetrachlorethane; the products were tetrachloroethylene, trichloroethylene, and 1,1-dichloroethylene, respectively. In vivo reaction rates were determined. No reaction was observed with 1,1,2,2-tetrachloroethane or 1,1,1-trichloroethane. Purified cytochrome P-450CAM was used to measure dissociation constants of polychlorinated ethanes for the enzyme active site. Observed rates and dissociation constants were used to predict the course of a reaction with the three substrates simultaneously. Data obtained from experiments with P. putida G786 generally followed the simulated reaction curves. Oxygen suppressed the reductive dechlorination reactions and, in the case of 1,1,1,2-tetrachlorethane, 2,2,2-trichloroacetaldehyde was formed. Significant rates of reductive dechlorination were observed at 5% oxygen suggesting that these reactions could occur under partially aerobic conditions. These studies highlight the potential to use an aerobic bacterium, P. putida G786, under a range of oxygen tensions to reductively dehalogenate mixed wastes which are only degraded at very low rates by obligately anaerobic bacteria.

=> d his

(FILE 'HOME' ENTERED AT 11:43:23 ON 14 JAN 2003)

FILE 'MEDLINE, CAPLUS, BIOSIS, BIOTECHDS, SCISEARCH, EMBASE' ENTERED AT 11:44:23 ON 14 JAN 2003

```
7515 S MONOOXYGENASE AND OXIDATION
L1
           4386 DUP REM L1 (3129 DUPLICATES REMOVED)
L2
             28 S L2 AND CYTOCHROME P-450CAM
L3
              O S L3 AND HOLOGENATED AROMATIC
L4
L5
              0 S L3 AND HALOGENATED AROMATIC
              0 S L3 AND HALO AROMATIC
Ь6
L7
             5 S L3 AND AROMATIC
             0 S L3 AND HALOGENATED SUBSTRATE
Г8
             0 S L2 AND HALOGENATED AROMATIC SUBSTRATE
L9
             1 S L3 AND DICHLOROBENZENE
L10
             5 S L3 AND (BENZENE OR BIPHENYL)
L11
             1 S L3 AND (CHLORINE)
L12
=> log y
```

COST IN U.S. DOLLARS SINCE FILE TOTAL ENTRY SESSION FULL ESTIMATED COST 93.26 93.68 TOTAL DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) SINCE FILE ENTRY SESSION CA SUBSCRIBER PRICE -3.91 -3.91

STN INTERNATIONAL LOGOFF AT 12:03:16 ON 14 JAN 2003

End of Result Set Print Generate Collection

L6: Entry 1 of 1

File: USPT

Oct 30, 2001

DOCUMENT-IDENTIFIER: US 6309883 B1

TITLE: Methods and compositions for cellular and metabolic engineering

Detailed Description Text (81):
Some examples of chemical targets for bioremediation include but are not limited to benzene, xylene, and toluene, camphor, naphthalene, halogenated hydrocarbons, polychlorinated biphenyls (PCBs), trichlorethylene, pesticides such as pentachlorophenyls (PCPs), and herbicides such as atrazine.

CLAIMS:

- 74. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid encodes one or more enzyme selected from the group consisting of: an acylase, a dioxygenase, a monooxygenase, a carotenoid synthetic enzyme, a hydrolytic enzyme, a catabolic enzyme, a dibenzothiopene catabolizing enzyme, a nitroreductase, a benzene degrading enzyme, a nitrobenzene degrading enzyme, a nitrotoluene degrading enzyme, a toxin degrading enzyme, an industrial chemical degrading enzyme, an herbicide degrading enzyme, a cellulose degrading enzyme, a pesticide degrading enzyme a pollutant degrading enzyme, a xylene degrading enzyme a toluene degrading enzyme, a camphor degrading enzyme, a naphthalene degrading enzyme, a halogenated hydrocarbon degrading enzyme, a biphenyl degrading enzyme, a polychlorinated biphenyl (PCB) degrading enzyme, a polycyclic aromatic hydrocarbon (PHA) degrading enzyme, a polyhydroxybutyrate (PHB) degrading enzyme, a trichlorethylene degrading enzyme, a pentachlorophenyl (PCP) degrading enzyme, a trichloroethylene degrading enzyme, a paranitrobenzyl, esterase, a sesquiterpene synthase, an expandase, a penicillin amidase, a penicillin G amidase, an enzyme which modifies 7-aminodeacetooxycephalosporanic acid (7-ADCA), an enzyme which modifies a semi-synthetically produced cephalosporin, and an enzyme which modifies penicillin V.
- 82. The method of claim 81, wherein the one or more toxin, industrial chemical, herbicide or pollutant comprises one or more of: benzene, xylene, toluene, camphor, naphthalene, a halogenated hydrocarbon, a polychlorinated biphenyl (PCB), a polycyclic aromatic hydrocarbon (PHA), a trichlorethylene, a pentachlorophenyl (PCP) or trichloroethylene.
- 103. The method of claim 102, the one or more toxin, industrial chemical, herbicide or pollutant comprising one or more of: benzene, xylene, toluene, camphor, naphthalene, a halogenated hydrocarbon, a polychlorinated biphenyl (PCB), a polycyclic aromatic hydrocarbon (PHA), a trichlorethylene, a pentachlorophenyl (PCP) or trichloroethylene.
- 126. The method of claim 1 or 3, wherein the screening comprises monitoring one or more enzymatic activities of one or more enzymes selected from the group consisting of: an acylase, a dioxygenase, a monooxygenase, a carotenoid synthetic enzyme, a hydrolytic enzyme, a catabolic enzyme, a nitroreductase, a benzene degrading enzyme, a nitrobenzene degrading enzyme, a nitrotoluene degrading enzyme, a toxin degrading enzyme, an industrial chemical degrading enzyme, an herbicide degrading enzyme, a cellulose degrading enzyme, a pesticide degrading enzyme a pollutant degrading enzyme, a xylene degrading enzyme, a toluene degrading enzyme, a camphor degrading enzyme, a naphthalene degrading enzyme, a halogenated hydrocarbon degrading enzyme, a polychlorinated biphenyl (PCB) degrading enzyme, a polycyclic aromatic hydrocarbon (PHA) degrading enzyme, a polyhydroxybutyrate (PHP) degrading enzyme, a trichlorethylene degrading enzyme, a pentachlorophenyl (PCP) degrading enzyme, a

trichloroethylene degrading enzyme, a paranitrobenzyl, esterase, a sesquiterpene synthase, an expandase, a penicillin amidase, a penicillin G amidase, an enzyme which modifies 7-aminodeacetooxycephalosporanic acid (7-ADCA), an enzyme which modifies a semi-synthetically produced cephalosporin, and an enzyme which modifies penicillin V.

128. The method of claim 127 wherein the one or more toxin, industrial chemical, herbicide or pollutant comprises one or more of: benzene, xylene, toluene, camphor, naphthalene, a halogenated hydrocarbon, a polychlorinated biphenyl (PCB), a polycyclic aromatic hydrocarbon (PHA), a trichlorethylene, a pentachlorophenyl (PCP) or trichloroethylene.

WEST

End of Result Set

Generate Collection Print

L6: Entry 1 of 1

File: USPT

Oct 30, 2001

US-PAT-NO: 6309883

DOCUMENT-IDENTIFIER: US 6309883 B1

TITLE: Methods and compositions for cellular and metabolic engineering

DATE-ISSUED: October 30, 2001

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Minshull; Jeremy San Francisco CA Stemmer; Willem P. C. Los Gatos CA

US-CL-CURRENT: 435/440; 435/6, 536/23.1, 536/24.3

CLAIMS:

What is claimed is:

1. A method of recombining one or more nucleic acids, the method comprising:

introducing one or more sets of nucleic acids into a plurality of cells, thereby providing a plurality of modified cells, each of the plurality of modified cells comprising at least one member of the one or more sets of nucleic acids;

transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells;

permitting recombination to occur between the at least one member of the one or more sets of nucleic acids and a nucleic acid present in the second of the plurality of modified cells to produce a recombinant nucleic acid;

introducing the recombinant nucleic acid into a third cell and permitting recombination between the recombinant nucleic acid and a third member present in a third cell of the plurality of modified cells, or between the recombinant nucleic acid and the first member or the second member, thereby producing a further recombined nucleic acid; and,

screening the further recombined nucleic acid for one or more properties or one or more encoded activities, thereby providing a selected recombinant nucleic acid.

- 2. The method of claim 1, comprising further recombining the selected recombinant nucleic acid with one or more additional nucleic acids and selecting the resulting further recombined nucleic acid to produce a further recombined selected nucleic acid.
- 3. A method of recombining one or more nucleic acids, the method comprising:

introducing one or more sets of nucleic acids into a plurality of cells, thereby providing a plurality of modified cells, each of the plurality of modified cells comprising at least one member of the one or more sets of nucleic acids;

transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells; permitting recombination to occur between the first member and a second member present in the second of the plurality of modified cells, thereby producing a recombinant nucleic acid;

screening the recombinant nucleic acid for one or more properties or one or more encoded activities; and,

further recombining the selected recombinant nucleic acid with one or more additional nucleic acid, or with the first or second nucleic acid, thereby producing a further recombined selected nucleic acid.

- 4. The method of claim 2 or 3, comprising screening the further recombined selected nucleic acid for one or more encoded activities, thereby providing a multiply recombined multiply selected nucleic acid.
- 5. The method of claim 2 or 3, wherein the further recombining comprises in vitro recombination.
- 6. The method of claim 5, wherein the further recombining comprises recursive in vitro recombination.
- 7. The method of claim 2 or 3, wherein the further recombining comprises in vivo recombination.
- 8. The method of claim 7, wherein the further recombining step comprises recursive in vivo recombination.
- 9. The method of claim 1 or 3, wherein the one or more sets of nucleic acids comprise one or more nucleic acid produced by in vitro sequence recombination.
- 10. The method of claim 1 or 3, wherein the one or more sets of nucleic acids comprise one or more nucleic acid produced by recursive in vitro recombination.
- 11. The method of claim 1 or 3, wherein the one or more sets of nucleic acids comprise one or more nucleic acid produced by in vivo recombination.
- 12. The method of claim 1 or 3, wherein the one or more sets of nucleic acids comprise one or more nucleic acid produced by recursive in vivo sequence recombination.
- 13. The method of claim 1 or 3, wherein the one or more sets of nucleic acids comprise one or more nucleic acid produced by mutation.
- 14. The method of claim 13, wherein the one or more sets of nucleic acids are produced by error prone PCR.
- 15. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises packaging members of one or more of the one or more sets into phage vectors and transducing the resulting phage library into a plurality of cells, thereby producing the plurality of modified cells.
- 16. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises packaging members of one or more of the one or more sets into viral vectors and transducing the

resulting viral library into a plurality of cells, thereby producing the plurality of modified cells.

- 17. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises electroporating members of one or more of the one or more sets into a plurality of cells, thereby producing the plurality of modified cells.
- 18. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises electronic pulse introduction of members of one or more of the one or more sets into a plurality of cells, thereby producing the plurality of modified cells.
- 19. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises biolistically introducing members of one or more of the one or more sets into a plurality of cells, thereby producing the plurality of modified cells.
- 20. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises transferring members of one or more of the one or more sets into a plurality of cells via conjugative transfer, thereby producing the plurality of modified cells.
- 21. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises transferring one or more of the one or more sets into the plurality of cells by fusing one or more cells comprising one or more members of the one or more sets with a plurality of cells, thereby producing the plurality of modified cells.
- 22. The method of claim 1 or 3, wherein the step of introducing the one or more sets of nucleic acids into the plurality of cells comprises transferring one or more members of the one or more sets of nucleic acids into a plurality of cells by fusing one or more library cells comprising members of the one or more sets with the one or more of the plurality of cells, wherein the fusing is induced by incubation of the library cells or the plurality of cells, or both, with a viral protein, or a chenmical agent.
- 23. The method of claim 22, wherein the viral protein comprises one or more of: an influenza protein, an influenza viral hemagglutinin protein, HSV-1 g B, or HSV-1 g D.
- 24. The method of claim 22, wherein the chemical agent is PEG.
- 25. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells comprises packaging at least one member of one or more of the one or more sets into at least one phage vector and transducing the resulting at least one phage vector into the second modified cell.
- 26. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells comprises packaging at least one member of one or more of the one or more sets into at least one viral vector and transducing the resulting at least one viral vector into the second modified cell.
- 27. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells comprises electroporating at least one member of one or more of the one or more sets into the second modified cell.

- 28. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells comprises electronic pulse transfer of at least one member of one or more of the one or more sets into the second modified cell.
- 29. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells comprises biolistically transferring at least one member of one or more of the one or more sets into the second modified cell.
- 30. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells is performed via conjugative transfer of the first member from the first modified cell into the second modified cell.
- 31. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells is performed by fusing the first and second cell.
- 32. The method of claim 1 or 3, wherein the step of transferring at least a first member of the one or more sets of nucleic acids from a first of the plurality of modified cells into at least a second of the plurality of modified cells is performed by fusing the first and second cell, wherein the fusing is induced by incubation of the first and second cells with a viral protein, or a chemical agent.
- 33. The method of claim 32, wherein the viral protein comprises one or more of: an influenza protein, an influenza viral hemagglutinin protein, HSV-1 g B, or HSV-1 g D.
- 34. The method of claim 32, wherein the chemical agent is PEG.
- 35. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell comprises packaging the recombinant nucleic acid into at least one phage vector and transducing the resulting at least one phage vector into the third cell.
- 36. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell comprises packaging the recombinant nucleic acid into at least one viral vector and transducing the resulting at least one viral vector into the third cell.
- 37. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acids into the third cell comprises electroporating the recombinant nucleic acid into the third cell.
- 38. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell comprises pulse introducing the recombinant nucleic acid into the third cell.
- 39. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell comprises biolistically introducing the recombinant nucleic acid into the third cell.
- 40. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell is performed via conjugative transfer of the recombinant nucleic acid into the third cell.

4 of 12 1/14/03 1:14 PM

- 41. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell comprises fusing the second and third cells.
- 42. The method of claim 1 or 3, wherein the step of introducing the recombinant nucleic acid into the third cell comprises fusing the second and third cells, wherein the fusing is induced bit incubation of the second and third cells with a viral protein, or a chemical agent.
- 43. The method of claim 42, wherein the viral protein comprises one or more of: an influenza protein, an influenza viral hemagglutinin protein, HSV-1 g B, or HSV-1 g D.
- 44. The method of claim 42, wherein the chemical agent is PEG.
- 45. The method of claim 1 or 3, wherein the plurality of modified cells comprise one or more mutator cells.
- 46. The method of claim 45, wherein the mutator cells are selected from the group consisting of: Mut L cells, Mut S cells, Mut D cells, Mut T cells, Mut H cells, and Human Ataxia Telengiecta cells.
- 47. The method of claim 1 or 3, wherein a plurality of members of the one or more sets of nucleic acids are at least about 50% identical.
- 48. The method of claim 1 or 3, wherein the members of the one or more sets of nucleic acids are at least about 70% identical.
- 49. The method of claim 1 or 3, wherein the members of the one or more sets of nucleic acids are at least about 80% identical.
- 50. The method of claim 1 or 3, wherein the members of the one or more sets of nucleic acids are at least about 90% identical.
- 51. The method of claim 1 or 3, wherein the members of the one or more sets of nucleic acids differ from each other in about 5 to about 20 positions.
- 52. The method of claim 1 or 3, wherein at least one of the one or more sets of nucleic acids have less than 10 members.
- 53. The method of claim 1 or 3, wherein at least one of the one or more sets of nucleic acids have more than 10.sup.5 members.
- 54. The method of claim 1 or 3, wherein at least one of the one or more sets of nucleic acids have more than 10.sup.7 members.
- 55. The method of claim 1 or 3, wherein at least one of the one or more sets of nucleic acids have more than 10.sup.9 members.
- 56. The method of claim 1 or 3, wherein at least one member of the one or more sets of nucleic acids is a full-length gene.
- 57. The method of claim 1 or 3, wherein at least one member of the one or more sets of nucleic acids is cloned into a vector which supplies one or more of: a promoter, a polyadenylation sequence, or a regulatory sequence.
- 58. The method of claim 1 or 3, wherein the members of the one or more sets of nucleic acids are allelic or species variants.
- 59. The method of claim 1 or 3, wherein at least one member of the plurality of modified cells is selected or derived from one or more of: a bacterial cell, gram-negative cell, gram-positive cell, a Streptomycetes cell, an Actinomycetes cell, a Corynebacteria cell, a Penicillium cell, a Bacillus cell, an

Escherichia coli cell, a Pseudomonas cell, a Salmonella cell, an Erwinia cell, a eukaryotic cell, a mammalian cell, a mouse cell, a hamster cell, a primate cell, a human cell, an established cell line cell, a primary cell culture cell, a stem cell, an embryonic stem cell, a zygotes cell, a fibroblast cell, a lymphocyte cell, a Chinese hamster ovary (CHO) cell, a mouse fibroblast cell, an NIH3T3 cell, a kidney cell, a liver cell, a muscle cell, a skin cell, a plant cell, a maize cell, a rice cell, a wheat cell, a cotton cell, a soybean cell, a sugarcane cell, a tobacco cell, an arabidopsis cell; a fish cell, an algal cell, a fungal cell, a Penicillium cell, a Fusarium cell, an Aspergillus cell, a Podospora cell, a Neurospora cell, an insect cell, a yeast cell, a Picchia cell, a Saccharomyces cell, or a nitrogen-fixation symbiotic cell.

- 60. The method of claim 1 or 3, wherein at least one member of the plurality of modified cells is selected or derived from a tissue or organism selected from the group consisting of: a plant, a bacteria, a fungus, an algae, an intact animal tissue, a tissue culture, and an animal embryo.
- 61. The method of claim 1 or 3, wherein at least one member of the plurality of modified cells is selected or derived from one or more of: E. coli, lactobacilli, Streptomycetes, Actinomycetes or filamentous fungi.
- 62. The method of claim 1 or 3, wherein at least one member of the plurality of modified cells is selected for one or more of: pathogenicity, substrate range, environmental hardiness, presence of one or more key intermediates, ease of genetic manipulation, or likelihood of promiscuous transfer of genetic information to other organisms.
- 63. The method of claim 1 or 3, wherein at least one member of the plurality of modified cells is selected or derived from one or more cell which comprises a biphenyl catabolizing pathway.
- 64. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid comprises one or more of: a plasmid, a cosmid, a chromosome, an episome, a YAC, a phage, a filamentous phage, a phage P1 clone, or a viral vector.
- 65. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid comprises cleaved genomic DNA.
- 66. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid comprises amplified genomic DNA.
- 67. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid comprises one or more metabolic pathway nucleic acids which encode at least one metabolic pathway.
- 68. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid are provided in or selected from a library of nucleic acids selected from the group consisting of: a plasmid library, a cosmid library, a phage library, a chromosome library, a filamentous phage library, and a viral library.

- 69. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid are provided in or selected from a library of nucleic acids comprising variants of a single gene.
- 70. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid are provided in or selected from a library of nucleic acids comprising variants of more than one gene.
- 71. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid are provided in or selected from a library of nucleic acids comprising one or more genes in a biochemical pathway.
- 72. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid are provided in or selected from a library of genes isolated from one or more of: a bacteria, an Alcaligenes, a Zoogloea, a Rhizobium, a Bacillus, a Azobacter, or a eukaryote.
- 73. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid comprises a nucleic acid which encodes a regulatory gene.
- 74. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid encodes one or more enzyme selected from the group consisting of: an acylase, a dioxygenase, a monooxygenase, a carotenoid synthetic enzyme, a hydrolytic enzyme, a catabolic enzyme, a dibenzothiopene catabolizing enzyme, a nitroreductase, a benzene degrading enzyme, a nitrobenzene degrading enzyme, a nitrotoluene degrading enzyme, a toxin degrading enzyme, an industrial chemical degrading enzyme, an herbicide degrading enzyme, a cellulose degrading enzyme, a pesticide degrading enzyme a pollutant degrading enzyme, a xylene degrading enzyme a toluene degrading enzyme, a camphor degrading enzyme, a naphthalene degrading enzyme, a halogenated hydrocarbon degrading enzyme, a biphenyl degrading enzyme, a polychlorinated biphenyl (PCB) degrading enzyme, a polycyclic aromatic hydrocarbon (PHA) degrading enzyme, a polyhydroxybutyrate (PHB) degrading enzyme, a trichlorethylene degrading enzyme, a pentachlorophenyl (PCP) degrading enzyme, a trichloroethylene degrading enzyme, a paranitrobenzyl, esterase, a sesquiterpene synthase, an expandase, a penicillin amidase, a penicillin G amidase, an enzyme which modifies 7-aminodeacetooxycephalosporanic acid (7-ADCA), an enzyme which modifies a semi-synthetically produced cephalosporin, and an enzyme which modifies penicillin V.
- 75. The method of claim 74, wherein the enzyme is a polyhydroxybutyrate (PHB) degrading enzyme, wherein the one or more sets of nucleic acids are derived from one or more of: an Alcaligenes bacteria, a Zoogloea bacteria, a Rhizobium bacteria, a Bacillus bacteria, or an Azobacter bacteria.
- 76. The method of claim 74, wherein the enzyme is a a biphenyl degrading enzyme and wherein the enzyme is expressed in at least one host cell which comprises a biphenyl catabolizing pathway.

- 77. The method of claim 74, wherein the enzyme is a cellulose degrading enzyme and wherein the one or more sets of nucleic acids are derived from one or more Agrobacterium tumefaciens.
- 78. The method of claim 74, wherein the enzyme is a carotenoid synthetic enzyme and wherein the one or more sets of nucleic acids are derived from one or more of: Myxococcus xanthus, Rhodobacter sphaeroides, Thermus thermophilus, Erwina uredovora, Haematococcus pluvialis, E. coli, E. herbicola, or R. capsulatus.
- 79. The method of claim 1, 2, or 3, wherein one or more member of the one or more sets of nucleic acids, the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid encodes one or more enzyme which is resistant to inactivation by one or more epoxide.
- 80. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expres, sed in the organism, with a new or improved ability to convert a pollutant into a nutrient source.
- 81. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism, with a new or improved ability to degrade one or more toxin, industrial chemical, herbicide, pesticide or pollutant.
- 82. The method of claim 81, wherein the one or more toxin, industrial chemical, herbicide or pollutant comprises one or more of: benzene, xylene, toluene, camphor, naphthalene, a halogenated hydrocarbon, a polychlorinated biphenyl (PCB), a polycyclic aromatic hydrocarbon (PHA), a trichlorethylene, a pentachlorophenyl (PCP) or trichloroethylene.
- 83. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid encode an enzyme with an improved catalytic activity, a new catalytic activity, altered substrate recognition, thermostability, stability in a non-aqueous solvent, or an altered expression level.
- 84. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism, with a new or improved resistance to the presence of one or more heavy metal.
- 85. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism one or more property selected from the group consisting of: modified growth rate, ability to secrete a desired compound, an ability to tolerate an increased temperature, and an ability erate one or more environmental stress.
- 86. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism, with a new or improved ability to reduce an organo-nitro compound or to permit the organism to survive in the presence of an organo-nitro compound.


- 87. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism, with new or improved utilization of a nutrient source.
- 88. The method of claim 87, wherein the nutrient source is selected from the group consisting of: lactose, whey, galactose, mannitol, xylan, cellobiose, cellulose and sucrose.
- 89. The method of claim 87, wherein the improved utilization of a nutrient source provides for production of compounds selected from the group consisting of: ethanol, tryptophan, a rhamnolipid surfactant, xanthan gum, polysaccharide xanthan gum and polyhydroxylalkanoate.
- 90. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism, new or improved production of one or more product selected from the group consisting of: ethanol, tryptophan, a rhamnolipid surfactant, xanthan gum, polysaccharide xanthan gum, polyhydroxylalkanoate, phenylalanine, and 2-keto-L-gluconic acid.
- 91. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid provides one or more organism, when expressed in the organism, with a new or improved ability to produce one or more metabolic intermediate.
- 92. The method of claim 91, wherein the metabolic intermediate is selected from the group consisting of: an antibiotic, a vitamin, an amino acid, phenylalanine, an aromatic amino acid, ethanol, butanol, polysaccharide xanthan gum, xanthan gum, bacterial cellulose, a peptide, and a lipid.
- 93. The method of claim 1, 2, or 3, wherein the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid encodes an enzyme which produces one or more compound selected from the group consisting of: a polyketide, a dye, a vitamin, an antibiotic, a carotenoid, a terpenoid, and an isoprenoid.
- 94. The method of claim 93, wherein the dye is indigo.
- 95. The method of claim 93, wherein the vitamin is vitamin C.
- 96. The method of claim 93, wherein the antibiotic is selected from the group consisting of: a peptide, a peptidolactone, a thiopeptide, a beta-lactam, a glycopeptide, a lantibiotic, a microcin, a polyketide-derived antibiotic, an anthracyclin, a tetracyclin, a macrolide, an avermectin, a polyether, an ansamycins, chloramphenicol, an aminoglycoside, an aminocyclitol, a polyoxin, an agrocin, mederrhodin, dihydrogranatirhodin, 6-deoxyerythromycin A, isovalerylspiramycin, a hybrid macrolide and an isoprenoid.
- 97. The method of claim 93, wherein the polyketide is an antibiotic.
- 98. The method of claim 93, wherein the polyketide is selected from the group consisting of: tetracycline, erythromycin, an anti-cancer agent, daunomycin, an immunosuppressant, FK506, rapamycin, monesin and avermectin.
- 99. The method of claim 93, wherein the isoprenoid is selected from the group consisting of: an antibacterial isoprenoid and an antifungal isoprenoid.

9 of 12 1/14/03 1:14 PM

- 100. The method of claim 93, wherein the carotinoid is selected from the group consisting of: a ketocarotenoid, a myxobacton, a spheroidene, a spheroidenone, a lutein, an astaxanthin, a violaxanthin, a 4-ketorulene, a myxoxanthrophyll, an echinenone, a lycopene, a zeaxanthin, a monoglucoside, a diglucoside, an alpha carotene, a beta carotene, a gamma carotene, a delta carotene, a cryptoxanthin monoglucoside and a neoxanthin.
- 101. The method of claim 1 or 3, further comprising propagating the first, second, or third cell, in culture.
- 102. The method of claim 1 or 3, wherein the screening comprises monitoring bioremediation or biodegradation of one or more toxin, industrial chemical, herbicide, pesticide or pollutant.
- 103. The method of claim 102, the one or more toxin, industrial chemical, herbicide or pollutant comprising one or more of: benzene, xylene, toluene, camphor, naphthalene, a halogenated hydrocarbon, a polychlorinated biphenyl (PCB), a polycyclic aromatic hydrocarbon (PHA), a trichlorethylene, a pentachlorophenyl (PCP) or trichloroethylene.
- 104. The method of claim 1 or 3, wherein the screening step is performed in the same cell type as the recombinant cell is produced in.
- 105. The method of claim 1 or 3, wherein the screening step is performed in a different cell type than the recombinant cell is produced in.
- 106. The method of claim 1 or 3, wherein the screening comprises monitoring one or more reporter gene selected from the group consisting of: luciferase, green fluorescence protein, and .beta.-galactosidase.
- 107. The method of claim 1 or 3, wherein the screening comprises monitoring one or more of: fluorescence, bioluminescence, colony size, cell growth rate, a chromogenic substrate, or halo formation.
- 108. The method of claim 1 or 3, wherein the screening comprises performing an ELISA assay.
- 109. The method of claim 1 or 3, wherein the screening comprises performing a cell-cell activity assay.
- 110. The method of claim 2 or 3, wherein the screening comprises monitoring differential expression of a protein or nucleic acid expressed in a screened cell comprising the recombinant nucleic acid, the further recombined nucleic acid, or the further recombined selected nucleic acid.
- 111. The method of claim 1 or 3, wherein the screening comprises performing FACS.
- 112. The method of claim 1 or 3, wherein the screening comprises performing two-color FACS.
- 113. The method of claim 1 or 3, wherein the screening comprises monitoring gel microdroplets.
- 114. The method of claim 1 or 3, wherein the screening comprises detecting one or more molecule by mass spectometry.
- 115. The method of claim 1, 2, or 3, wherein a selected cell comprising the recombinant nucleic acid, the further recombined nucleic acid, the further recombined selected nucleic acid or the multiply recombined multiply selected nucleic acid is selected in a chemostat.

- 116. The method of claim 1 or 3, wherein the screening comprises selecting for one or more of: an improved catalytic activity, a new catalytic activity, altered substrate recognition, thermostability, stability in a non-aqueous solvent, or an altered expression level.
- 117. The method of claim 1 or 3, wherein the screening comprises selecting one or more organism comprising the recombinant nucleic acid for one or more property selected from the group consisting of: a modified growth rate, an ability to secrete a desired compound, an ability to tolerate an increased temperature, and an ability to tolerate one or more environmental stresses.
- 118. The method of claim 1 or 3, wherein the screening comprises monitoring the presence or absence of one or more secondary metabolite selected from the group consisting of: a polyketide, a dye, a vitamin, an antibiotic, a carotenoid, a terpenoid, and an isoprenoid.
- 119. The method of claim 118, wherein the dye is indigo.
- 120. The method of claim 118, wherein the vitamin is vitamin C.
- 121. The method of claim 118, wherein the antibiotic is selected from the group consisting of: a peptide, a peptidolactone, a thiopeptide, a beta-lactam, a glycopeptide, a lantibiotic, a microcin, a polyketide-derived antibiotic, an anthracyclin, a tetracyclin, a macrolide, an avermectin, a polyether, an ansamycins, chloramphenicol, an aminoglycoside, an aminocyclitol, a polyoxin, an agrocin, mederrhodin, dihydrogranatirhodin, 6-deoxyerythromycin A, isovalerylspiramycin, a hybrid macrolide and an isoprenoid.
- 122. The method of claim 118, wherein the polyketide is an antibiotic.
- 123. The method of claim 118, wherein the polyketide is selected from the group consisting of: tetracycline, erythromycin, an anti-cancer agent, daunomycin, an immunosuppressant, FK506, rapamycin, monesin and avermectin.
- 124. The method of claim 118, wherein the isoprenoid is selected from the group consisting of: an antibacterial isoprenoid and an antifungal isoprenoid.
- 125. The method of claim 118, wherein the carotinoid is selected from the group consisting of: a ketocarotenoid, a myxobacton, a spheroidene, a spheroidenone, a lutein, an astaxanthin, a violaxanthin, a 4-ketorulene, a myxoxanthrophyll, an echinenone, a lycopene, a zeaxanthin, a monoglucoside, a diglucoside, an alpha carotene, a beta carotene, a gamma carotene, a delta carotene, a cryptoxanthin monoglucoside and a neoxanthin.
- 126. The method of claim 1 or 3, wherein the screening comprises monitoring one or more enzymatic activities of one or more enzymes selected from the group consisting of: an acylase, a dioxygenase, a monooxygenase, a carotenoid synthetic enzyme, a hydrolytic enzyme, a catabolic enzyme, a nitroreductase, a benzene degrading enzyme, a nitrobenzene degrading enzyme, a nitrotoluene degrading enzyme, a toxin degrading enzyme, an industrial chemical degrading enzyme, an herbicide degrading enzyme, a cellulose degrading enzyme, a pesticide degrading enzyme a pollutant degrading enzyme, a xylene degrading enzyme, a toluene degrading enzyme, a camphor degrading enzyme, a naphthalene degrading enzyme, a halogenated hydrocarbon degrading enzyme, a polychlorinated biphenyl (PCB) degrading enzyme, a polycyclic aromatic hydrocarbon (PHA) degrading enzyme, a polyhydroxybutyrate (PHP) degrading enzyme, a trichlorethylene degrading enzyme, a pentachlorophenyl (PCP) degrading enzyme, a trichloroethylene degrading enzyme, a paranitrobenzyl, esterase, a sesquiterpene synthase, an expandase, a penicillin amidase, a penicillin G amidase, an enzyme which modifies 7-aminodeacetooxycephalosporanic acid (7-ADCA), an enzyme which modifies a semi-synthetically produced cephalosporin, and an enzyme which modifies penicillin V.

- 127. The method of claim 1 or 3, wherein the screening comprises monitoring degradation of one or more of: a toxin, an industrial chemical, an herbicide, a pesticide a pollutant, PHB, or cellulose.
- 128. The method of claim 127 wherein the one or more toxin, industrial chemical, herbicide or pollutant comprises one or more of: benzene, xylene, toluene, camphor, naphthalene, a halogenated hydrocarbon, a polychlorinated biphenyl (PCB), a polycyclic aromatic hydrocarbon (PHA), a trichlorethylene, a pentachlorophenyl (PCP) or trichloroethylene.
- 129. The method of claim 1 or 3, wherein the screening comprises monitoring synthesis of one or more carotenoid.
- 130. The method of claim 1 or 3, wherein the screening comprises monitoring resistance of an enzyme to an epoxide.
- 131. The method of claim 1 or 3, wherein the screening comprises monitoring resistance of a cell modified with the recombinant nucleic acid to a heavy metal.
- 132. The method of claim 1 or 3, wherein the screening comprises selecting an organism which expresses the recombinant nucleic acid for an ability to survive in the presence of an organo-nitro compound.
- 133. The method of claim 1 or 3, wherein the screening comprises selecting an organism for an ability to metabolize lactose, whey, galactose, mannitol, xylan, cellobiose, cellulose or sucrose.
- 134. The method of claim 1 or 3, wherein the screening comprises selecting, an organism for an ability to produce ethanol, tryptophan, a rhamnolipid surfactant, xanthan gum, polysaccharide xanthan gum, polyhydroxylalkanoate, phenylalanine, or 2-keto-L-gluconic acid.



L5: Entry 1 of 7 File: USPT Apr 23, 2002

US-PAT-NO: 6376254

DOCUMENT-IDENTIFIER: US 6376254 B1

TITLE: Biomimetic reagent system and its use

DATE-ISSUED: April 23, 2002

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY Bather; Wolfgang Lubeck DE

Duchstein; Hans-Jurgen Pinneberg DE
Hoffmann; Susanne Buchholz DE

ASSIGNEE-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY TYPE CODE

Drager Sicherheitstechnik GmbH DE 03

APPL-NO: 09/ 394969 [PALM]
DATE FILED: September 10, 1999

FOREIGN-APPL-PRIORITY-DATA:

COUNTRY APPL-NO APPL-DATE

DE 199 12 380 March 19, 1999

INT-CL: [07] G01 N 21/78

US-CL-ISSUED: 436/140; 436/167, 422/86, 422/88 US-CL-CURRENT: 436/140; 422/86, 422/88, 436/167

FIELD-OF-SEARCH: 436/140, 436/164, 436/167, 436/169, 436/181, 422/55, 422/86, 422/87,

422/88, 422/91

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

Search Selected Search ALL

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
4864041	September 1989	Hill	
5063164	November 1991	Goldstein	
5516636	May 1996	McCapra	
5712402	January 1998	Pinnavaia et al.	
6020288	February 2000	Nonomura et al.	
6028025	February 2000	Ying et al.	
6103892	August 2000	Breslow et al.	

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
3407 686	July 1985	DE	
39 02 402	June 1990	DE	

OTHER PUBLICATIONS

A.M. Gonsalves et al "State of the Art in the Development of Biomimetic Oxidation Catalysts" Journan of Molecular Catalysis A: Chemical, Vol. 113 (1996), pp. 209-211.

ART-UNIT: 1743

PRIMARY-EXAMINER: Snay; Jeffrey

ABSTRACT:

A biomimetic reagent system is provided containing an oxygen donor and a catalyst based on porphyrin, which are applied to a carrier. A device that contains the system is also provided for determining components of gas or vapor samples, especially aromatics, such as benzene. A process for hydroxylating aromatics, such as benzene, using the biomimetic reagent system is also provided.

30 Claims, 1 Drawing figures

isplay Form	. v	ysiwyg://26/h	tp://westbrs:80	02/bin/gatge=	&p_Message=&p_do	ccnt=1&p_d
		WES		www.	on anarassis and anarassis	ss::::::
End of Result Set	<u> </u>	nerate Colle	ection	Print		
L6: Entry 1 of 1		File	: USPT		Oct 3	30, 2001
US-PAT-NO: 6309883 DOCUMENT-IDENTIFIER: US	G 6309883 B1					
TITLE: Methods and comp	positions for ce	ellular a	nd metabo	lic engin	eering	
DATE-ISSUED: October 30), 2001					
INVENTOR-INFORMATION:	GT TIV		QM.	mn 275	CODE COUNT	IDV
NAME Minshull; Jeremy	CITY San Fi	rancisco	STA CA	TE ŽIP (LODE COUNT	K1
Stemmer; Willem P. C.	Los Ga		CA			
ASSIGNEE-INFORMATION:						
NAME CIT			ZIP CODE	COUNTR		E
Maxygen, Inc. Rec	lwood City	CA			02	
APPL-NO: 09/ 490642 DATE FILED: January 24	[PALM] 2000					
PARENT-CASE: This application is a C Ser. No. 08/650,400 fil Ser. No. 08/198,431 fil	led May 20, 1996	s, now U.	S. Pat. N	lo. 5,837,4	458; which is	a CIP

PAR

of Thi of Ser Ser No. 08/621,859 filed Mar. 25, 1996, now U.S. Pat. No. 6,117,679; and a CIP of Ser. No. 08/621,430 filed Mar. 25, 1996, now abandoned; and a CIP of Ser. No. 08/537,874 filed Mar. 4, 1996, now U.S. Pat. No. 5,830,721; which is the national phase of PCT/US95/02126 filed Feb. 17, 1995; and a CIP of Ser. No. 08/425,684 filed Apr. 18, 1995, now U.S. Pat. No. 5,834,252.

INT-CL: [07] $\underline{C12}$ \underline{N} $\underline{15/00}$, $\underline{C12}$ \underline{Q} $\underline{1/68}$, $\underline{C07}$ \underline{H} $\underline{21/02}$, $\underline{C07}$ \underline{H} $\underline{21/04}$

US-CL-ISSUED: 435/440; 435/6, 536/23.1, 536/24.3, 935/76, 935/77, 935/78

US-CL-CURRENT: 435/440; 435/6, 536/23.1, 536/24.3

FIELD-OF-SEARCH: 435/440, 435/6, 536/23.1, 536/24.3, 935/76, 935/77, 935/78

Search Selected

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

Search ALL

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
4683202	July 1987	Mullis	
4800159	January 1989	Mullis et al.	
4965188	October 1990	Mullis et al.	
4994368	February 1991	Goodman et al.	

4994379	February 1991	Chang	
5023171	June 1991	Ho et al.	
5032514	July 1991	Anderson et al.	435/138
5043272	August 1991	Hartley	
5093257	March 1992	Gray	435/202
5176995	January 1993	Sninsky et al.	
5187083	February 1993	Mullis	
5223408	June 1993	Goeddel et al.	435/69.3
5234824	August 1993	Mullis	
5279952	January 1994	Wu	435/172.3
5316935	May 1994	Arnold et al.	
5356801	October 1994	Rambosek et al.	435/195
5360728	November 1994	Prasher	435/189
5418149	May 1995	Gelfland et al.	
5422266	June 1995	Cormier et al.	
5489523	February 1996	Mathur	435/194
5502167	March 1996	Waldmann et al.	
5512463	April 1996	Stemmer	
5514588	May 1996	Stemmer	
5521077	May 1996	Khosla et al.	435/172.3
5541309	July 1996	Prasher	536/23.2
5556750	September 1996	Modrich et al.	
5556772	September 1996	Sorge et al.	
5605793	February 1997	Stemmer	435/6
5629179	May 1997	Mierendorf et al.	
5652116	July 1997	Grandi et al.	435/69.1
5679522	October 1997	Modrich et al.	
5714316	February 1998	Weiner et al.	
5723323	March 1998	Kauffman et al.	
5756316	May 1998	Schellenberger	
5763192	June 1998	Kauffman et al.	
5773267	September 1998	Jacobs	
5783431	July 1998	Peterson et al.	
5789228	August 1998	Lam et al.	
5795747	August 1998	Henco et al.	
5811238	September 1998	Stemmer et al.	
5814473	September 1998	Warren et al.	

2 of 13

5817483	October 1998	Kauffman et al.
5824469	October 1998	Horwitz et al.
5824485	October 1998	Thompson et al.
<u>5824514</u>	October 1998	Kauffman et al.
<u>5830696</u>	November 1998	Short
<u>5830721</u>	November 1998	Stemmer et al.
5834252	November 1998	Stemmer et al.
5837458	November 1998	Minshull et al.
5851813	December 1998	Desrosiers
5858725	January 1999	Crowe et al.
5876997	March 1999	Kretz
5925749	July 1999	Mathur et al.
5928905	July 1999	Stemmer et al.
5939250	August 1999	Short
5939300	August 1999	Robertson et al.
5942430	August 1999	Robertson et al.
5948666	September 1999	Callen et al.
5958672	September 1999	Short
5958751	September 1999	Murphy et al.
5962258	October 1999	Mathur et al.
5962283	October 1999	Warren et al.
5965408	October 1999	Short
5976862	November 1999	Kauffman et al.

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
0 252666 B1	January 1988	EP	
0911396A2	April 1999	EP	
552 266	April 1999	EP	
0911396A3	May 1999	EP	
0934999A1	August 1999	EP	
WO 93/18141	September 1963	WO	
WO 90/07576	July 1990	WO	
WO 90/14430	November 1990	WO	
WO 91/01087	February 1991	WO	
WO 91/07506	May 1991	WO	
WO 91/06570	May 1991 .	WO	
WO 91/16427	October 1991	WO	
WO 91/15581	October 1991	WO	
WO 92/07075	April 1992	WO	
WO 92/18645	October 1992	WO	
WO 93/02191	February 1993	WO	
WO 93/06213	April 1993	WO	

1/14/03 1:16 PM

WO	93/11237	June 1993	WO
WO	93/15208	August 1993	WO
WO	93/16192	August 1993	WO
WO	93/25237	December 1993	WO
WO	94/03596	February 1994	WO
WO	94/09817	May 1994	WO
WO	94/13804	June 1994	WO
WO	95/17413	June 1995	WO
WO	95/22625	August 1995	WO
WOS	95/22625	August 1995	WO
WO	96/33207	October 1996	WO
WO	97/07205	February 1997	MO
WO	97/20078	June 1997	WO
WO	97/25410	July 1997	WO
WO	97/35966	October 1997	MO
WO	97/35957	October 1997	WO
WO	97/44361	November 1997	MO
WO	97/48794	December 1997	WO
WO	97/48717	December 1997	MO
WO	97/48416	December 1997	WO
WO	98/01581	January 1998	WO
WO	98/00826	January 1998	WO
WO	98/13487	April 1998	MO
WO	98/13485	April 1998	WO
WO	98/27230	June 1998	WO
WO	98/24799	June 1998	WO
MO	98/31837	July 1998	WO
WO	98/28416	July 1998	WO
WO	98/36080	August 1998	WO
	98/41653	September 1998	WO
	98/41622	September 1998	WO
	98/41623	September 1998	WO
	98/48034	October 1998	WO
	98/42832	October 1998	WO
	98/58085	December 1998	WO
	99/07837	February 1999	WO
	99/08539	February 1999	WO
	99/10472	March 1999	WO
	99/10539	March 1999	WO
	99/19518	April 1999	WO
	99/21979	May 1999	WO
	99/23107	May 1999	WO
	99/23236	May 1999	MO
	99/41402	August 1999	WO
	99/41383	August 1999	WO
	99/41369	August 1999	WO
	99/45154	September 1999	WO
	99/57128	November 1999	MO
WO	99/65927	December 1999	WO

OTHER PUBLICATIONS

Prasher, Trends in Genetics 11(8): 320-323 (Aug. 1995).*
PCR Primer "A laboratory Manual" Eds. Dieffenbach and Dveksler, Cold Spring Harbor

```
Laboratory Press, 1995, pp. 583-621.*
Kunkel, Proc. Natl. Acad. Sci. (USA) 82 : 488-492 (1992).*
Lewis et al., Nucleic Acids Research 18(12) : 3439-3443 (1990).*
Rice et al., Proc. Natl. Acad. Sci. (USA) 89 : 5467-5471 (1992).*
Ippolito et al., Proc. Natl. Acad. Sci. (USA) 92 : 5017-5021 (May 1995).*
Atreya et al., "Construction of in-frame chimeric plant genes by simplified PCR
strategies, " Plant Mol. Biol., 19:517-522 (1992).
Bock et al., "Selection of single-stranded DNA molecules that bind and inhibit human
thrombin, "Nature, 355:564-566 (Feb. 2, 1992).
Shuldiner, A.R. et al., (1989) "Hybrid DNA artifact from PCR of closely related target
sequences," Nuc. Acids Res. 17(11): 4409.
Sandhu, G.S. et al., (1992) "Dual Asymmetric PCR: One-Step Construction of Synthetic
Genes, Biotechniques 12(01): 14-16.
Biotransformations, Pathogenesis, and Evolving Biotechnology, Program and Abstracts,
Pseudomonas '89, American Society for Microbiology and The University of Illinois, Jul.
9-13, 1989, Abstract Nos. 11-21--11-25, p. 26.
Smith et al. (1991) "Losalized sex in bacteria," Nature, 349: 29-31.
Michael et al., (c. 1997) "Thermostable Ligase-Mediated Incorporation of Mutagenic
Oligonucleotides During PCR Amplifications," Methods in Molecular Biology, vol. 67, PCR
Cloning Protocals: From Molecular Colning to Genetic Engineering pp. 189-195.
Ner et al. (1988) "Laboratory Methods: A Simple and Efficient Procedure for Generating
Random Point Mutations and for Codon Replacements Using Mixed Oligodeoxynucleotides,"
DNA 7: 127-134.
Prodromou (1992) "Recursive PCR: a novel technique for total gene synthesis," Protein
Engineering 5(8): 827-829.
Carter, "Improved Oligonucleotide-Directed Mutagenesis Using M13 Vectors," Methods in
Enzymology, 154:382 (1987).
Klug et al. (1991) "Creating chimeric molecules by PCR directed homologous DNA
recombination." Nuc. Acids Res. 19(10): 2793.
Rouwendal et al., (1993) "Simultaneous Mutagenesis of Multiple Sites: Application of
the Ligase Chain Reaction Using PCR Products instead of Oligonucleotides."
Biotechniques 15(1): 68-70, 72-74, 76.
Christians, F.C. et al., (1999) Directed evolution of thymidine kinase for AZT
phosphorylation using DNA family shuffling. Nature Biotechnology 17:259-264.
Fullen et al., (1994) "Genetic Algorithms and Recursive Ensemble Mutagenesis in Prtoein
Engineering" Complexity Int'l 1994, printed from website
http://www.csu.edu.au/ci/voll/fuellen/REM.html.
Lowman, H.B. et al., (1993) "Affinity Maturation of Human Growth Hormone by Monovalent
Phage Display" J. Mol. Biol. 234:654-578.
Minshull, J., Stemmer, W.P.C. (1999) Protein evolution by molecular breeding. Current
Opinion in Chemical Biology 3:284-290.
Patten, P.A. et al., (1997) "Application of DNA Shuffling to Pharmaceuticals and
Vaccines. " Current Opinion in Biotechnology 8:724-733.
Rapley (1994) "Enhancing PCR Amplification and Sequenceing Using DNA-Binding Proteins"
Molecular Biotechnology 2:295-298.
Robles et al. (1993) "Hydropathy and Molar volume Constraints on Combinatorial mutants
of the Photosynthetic Reaction Center" J. Mol. Biol. 232:242-252.
Youvan et al. (1992) "Recursive Ensemble Mutagenesis: A Combinatorial Technique for
Protein Engineering" from Parallel Problem Solving from Nature, 2, Manner eds., pp.
Appleyard, V.C.L. et al., "Secondary metabolite production in filamentous fungidisplayed," Mol. Gen. Gent. 247:338-342 (1995).
Arkin et al., "An Algorithm for Protein Engineering: Simulations of Recursive Ensemble
Mutagenesis" Proc. Natl. Acad. Sci. USA 89:7811-7815 (1992).
Armstrong, G.A., "Eubacteria Show Their Tru Colors: Genetics of Carotenoid Pigment
Biosynthesis from Microbes to Plants, " J. Bacteriol. 176(16):4795-4802 (1994).
Arnold, F.H., "Protein engineering for unusual environments," Cr. Opin. Biotechnol.
4:450-455 (1993).
Backman, K. et al., "Genetic Engineering of Metabolic Pathways Applied to the
Production of Phenylalanine, "Ann. NY Acad. Sci. 589:16-24 (1990).
Bailey, J.E., "Toward a Science of Metabolic Engineering," Science 252:1668-1675
(1991).
Beaudry et al., "Directed Evolution of an RNA Enzyme" Science 257:635-641 (1992).
Berger et al., "Phoenix Mutagenesis: One-Step Reassembly of Multiply Cleaved Plasmids
With Mixtures of Mutant and Wild-Type Fragments" Anal. Biochem 214:571-579 see pp.
571-578 (1993).
Berkhout et al., "In Vivo Selection of Randomly Mutated Retroviral Genomes" Nucleic
Acids Research 21:5020-5023 No. 22 (1993).
Botella, J.A. et al., "A cluster of structural and regulatory genes for light-induced
```

```
carotenogenesis in Myxococcus xanthus, " Eur. J. Biochem. 233:238-248 (1995).
Boudrant, J., "Microbial processes for ascorbic acid biosynthesis: a review," Enzyme
Microb. Technol. 12:322-329 (1990).
Brenner, V. et al., "Genetic construction of PCB degraders," Biodegradation 5:359-377
(1994).
Bryant, C. et al., "Cloning, Nucleotide Sequence, and Expression of the Nitroreductase
Gene from Enterobacter cloacae, " J. Biol. Chem. 266(7):4126-4130 (1991).
Cadwell et al., "Randomization of Genes by PCR Mutagenesis" PCR Methods and
Applications 2:28-33 (1992).
Cameron, D.C. et al., "Cellular and Metabolic Engineering," Applied Biochem. Biotech.
38:105-140 (1993).
Chakrabarty, A.M., "Microbial Degradation of Toxic Chemicals: Evolutionary Insights and
Practical Considerations, "Am. Soc. Micro. Biol. News 62:130-137 (1996).
Chater, K.F., "The improving prospects for yield increase by genetic engineering in
antibiotic-producing streptomycetes, "Bio/Technology 8:115-121 (1990).
Chen, C.W. et al., "Cloning and Expression of a DNA sequence conferring cepamycin C
production, "Bio/Technology 6:1222-1224 (1988).
Chen, K. et al., "Tuning the activity of an enzyme for unusual environments: Sequential
random mutagenesis of subtilisin E for catalysis in dimethylformamide," Proc. Natl.
Acad. Sci. U.S.A. 90:5618-5622 (1993).
Calogero Sabina et al., "In Vivo Recombination and the Production of Hybrid Genes,"
Microbiology letters, 1992, vol. 97, pp. 41-44.
Caren Robert et al., "Efficient Sampling of Protein Sequence Space for Multiple
Mutants, Biotechnology, May 12, 1994, vol. 12, pp. 517-520.
Dacre, J.C., "Nonspecificity of the Gibbs Reaction," Anal. Chem. 43(4):589-591 (1971).
Denome, S.A. et al., "Dibenzothiophene and Naphthalene in Pseudomonas Strains: Complete
DNA Sequence of an Upper Naphthalene Catabolic Pathway, " J. Bacteriol.
175(21):6890-6901 (1993).
de Souza, M.L. et al., "Cloning, Characterization, and Expression of a Gene Region from
Pseudomonas sp. Strain ADP Involved in the Dechlorination of Atrazine, " Appl. Environ.
Micro. 61(9):3373-3378 (1995).
Delagrave Simon et al., "Recursive Ensemble Mutagenesis," Protein Engineering, 1993,
vol. 6, No. 3, pp. 327-331.
Eaton, R.W. et al., "Formation of Indigo and Related Compounds from Indolecarboxylic
Acids by Aromatic Acid-Degrading Bacteria: Chromogenic Reactions for Cloning Genes
Encoding Dioxygenases That Act on Aromatic Acids, J. Bacteriol. 177(23):6983-6988
(1995).
Erickson B.D. et al., "Enhanced Biodegradation of Polychlorinated Biphenyls after
Site-Directed Mutagenesis of a Biphenyl Dioxygenase Gene, "App. Environ. Micro.
59(11):3858-3862 (1993).
Evnin, L.B. et al., "Substrate specificity of trypsin inestigated by using a genetic
selection, " Proc. Natl. Acad. Sci. U.S.A. 87:6659-6663 (1990).
Fernandez-Lafuente R. et al., "Additional stabilization of penicillin G acylase-agarose
derivatives by controlled chemical modification with formaldehyde," Enzyme Microb.
Technol. 14:489-495 (1992).
Ford, T. et al., "Metal-microbe interactions," Bioextraction and Biodeterioration of
Metals, p. 1-23.
Furukawa K. et al., "Gene-specific transposon mutagenesis of the
biphenyl/polychlorinated biphenyl-degradation-controlling bph operon in soil bacteria,"
Gene 98:21-28 (1991).
Furukawa K. et al., Gene Components Responsible for Discrete Substrate Specificity in
the Metabolism of Biphenyl (bph Operon) and Toluene (tod Operon), J. Bact.
175(16):5224-5232 (1993).
Furukawa K. et al., "Efficient Degradation of Trichloroethylene by a Hybrid Aromatic
Ring Dioxygenase, "J. Bacteriol. 176(7):2121-2123 (1994).

Goguen, B. et al., "Clonogenic cytotoxicity testing by microdrop encapsulation," Nature
363:189-190 (1995).
Hart, H.E. et al., "Scintillation proximity assay (SPA) -- a new method of immunoassay,"
Mol. Immunol. 16:265~267 (1979).
Hassan, H.M. et al., "Intracellular Production of Superoxide Radical and of Hydrogen
Peroxide by Redox Active Compounds, "Arch. Bioch. Biop. 196(2):385-395 (1979).
Hayase N. et al., "Pseudomonas putida KF715 bphABCD Operon Encoding Biphenyl and
Polychlorinated Biphenyl Degradation: Cloning, Analysis, and Expression in Soil
Bacteria, " J. Bacteriol. 172(2):1160-1164 (1990).
Hermes et al., "Searching Sequence Space by Definably Random Mutagenesis: Improving the
Catalytic Potency of an Enzyme" Proc. Natl. Acad. Sci. USA 87:696-700 (1990).
Heim et al., "Wavelength Mutations and Posttranslational Autoxidation of Green
Fluorescent Protein, Proc. Natl. Acad. Sci. USA 501-12504 (1994).
Hofer, B. et al., "The biphenyl/polychlorinated biphenyl-degradation locus (bph) of
```

wysiwyg://26/http://westbrs:8002/bin/gat...ge=&p_Message=&p_doccnt=1&p_doc_1=PTFFRO Pseudomonas sp. LB400 encodes four additional metabolic enzymes, " Gene 144:9-16 (1994). Hopwood, D.A., "Antibiotics: opportunities for genetic manipulation," Phil. Trans. R. Soc. Lond. B324:549-562 (1989). Hoshino T. et al., "Molecular Cloning and Sequence Analysis of the crtB Gene of Thermus thermophilus HB27, an Extreme Thermophile Producing Carotenoid Pigments," Appl. Environ. Micro. 59(9):3150-3153 (1993). Hunter, C.N. et al., "Introduction of New Carotenoids into the Bacterial Photosynthetic Apparatus by Combining the Carotenoid Biosynthetic Pathways of Erwinia herbicola and Rhodobacter sphaeroides, J. Bacteriol. 176(12):3692-3697 (1994). Hutchinson, C.R., "Prospects for the Discovery of New (Hybrid) Antibiotics by Genetic Engineering of Antibiotic-Producing Bacteria," Med. Res. Rev. 8(4):557-567 (1988). Hutchinson, C.R. et al., "Drug Development through the Genetic Engineering of Antibiotic-Producing Microorganisms, "Ann. NY Acad. Sci. 656:78-93 (1991). Hutchinson, C.R., "Drug Synthesis by Genetically Engineered Microorganisms," Bio/Technology 12:375-308 (1994). Ji, G. et al., "Bacterial resistance mechanisms for heavy metals of environmental concern, " J. Ind. Micro. 14:61-75 (1995). Jones et al., "Recombinant Circle PCR and Recombination PCR for Site-Specific Mutagenesis Without PCR Product Purification, "BioTechniques 12(4): pp. 528-534 (1992). Kim et al., "Human Immunodeficiency Virus Reverse Transcriptase" The Journal of Biological Chemistry 271, No. 9 pp. 4872-4878 (1996). Kajiwara, S. et al., "Isolation and functional identification of a novel cDNA for astaxanthin biosynthesis from Haematococcus pluvialis, and astaxanthin synthesis in Escherichia coli, Plant Mol. Bio. 29:343-352 (1995).

Kato, F. et al., "Carotenoid synthesis in Streptomyces setonii ISP5395 is induced by the gene crtS, whose product is similar to a sigma factor," Mol. Gen. Genet. 247:387-390 (1995). Kellogg, S.T. et al., "Plasmid Assisted Molecular Breeding: New Technique for Enhanced Biodegradation of Persistent Toxic Chemicals, "Science 214:1133-1135 (1981). Kleinkauf, H. et al., "A nonribosomal system of peptide biosynthesis," Eur. J. Biochem. 236:335-351 (1996). Krawiec, S., "Bacterial desulfurization of thiophenes: screening techniques and some speculations regarding the biochemical and genetic bases, " Devl. Indust. Microbiol. 31:103-114 (1990). Lang, H.P. et al., "Complete DNA Sequence, Specific Tn5 Insertion Map, and Gene Assignment of the Carotenoid Biosynthesis Pathway of Rhodobacter sphaeroides," J. Bacteriol. 177(8):2064-2073 (1995). Leung et al., "A Method for Random Mutagenesis of a Defined DNA Segment Using a Modified Polymerase Chain Reaction" 1:11-15, see pp. 11-14 (1989). Liao, J.C., "Modelling and analysis of metabolic pathways," Curr. Opin. Biotech. 4:211-216 (1993). Mandelbaum, R.T. et al., "Isolation and Characterization of a Pseudomonas sp. That Mineralizes the s-Triazine Herbicide Atrazine, "Appl. Environ. Micro. 61(4):1451-1457

Marks James et al., "By-Passing Immunization: Building High Affinity Human Antibodies by Chain Shuffling" Bio/Technology 10:779-782 (1992).

Marrs, B., "Mobilization of the Genes for Photosynthesis from Rhodopseudomonas capsulata by a Promiscuous Plasmid, "J. Bacteriol. 146(3):1003-1012 (1981). Martin, L. et al., "Cloning and sequencing of the pac gene encoding he penicillin G acylase of Bacillus megaterium ATCC 14945," FEMS Microbiol. Lett. 125:287-292 (1995). Matthysse, A.G. et al., "Genes Requiered for Cellulose Synthesis in Agrobacterium tumefaciens," J. Bacteriol. 177(4):1069-1075 (1995).

McDaniel, R. et al., "Engineered Biosynthesis of Novel Polyketides," Science 262:1546-1550 (1995).

Meyerhans Andreas et al., "DNA Recombination during PCR," Nucleic Acids Research, 1990, vol. 18, No. 7, pp. 1687-1691. Misawa, N. et al., "Structure and Functional Analysis of a Marine Bacterial Carotenoid

biosynthesis Gene Cluster and Astaxanthin Biosynthetic Pathway Proposed at the Gene Level, " J. Bacteriol. 177(22):6576-6584 (1995).

Moore, J.C. et al., "Directed evolution of a para-nitrobenzyl esterase for aqueous-organic solvents, "Nature Biotechnology 14:458-467 (1996). Murdock, D. et al., "Construction of Metabolic Operons Catalyzing the De Novo Biosynthesis of Indigo in Escherichia coli, Bio/Technology 11:381-386 (1993).

Nisbet, L.J. et al., "Developments in Microbial Products Screening," Ann. Rep. Med. Chem. 21:149-157 (1986).

Oliphant et al., "Cloning of Random-Sequence Oligodeoxynucleotides" Gene 44:177-183 (1988).

```
Omura, S., "Philosophy of new Drug Discovery," Microbio. Rev. 50(3):259-279 (1986).
Paau, A.S., "Improvement of Rhizogium Inoculants by Mutation, Genetic Engineering and
Formulation, " Biotech. Adv. 9:173-184 (1991).
Piepersberg, W., Pathway Engineering in Secondary Metabolite-Producing Actinomycetes,
Crit. Rev. Biotechnol. 14:251-285 (1994).
Poirier, Y. et al., "Polyhydroxybutyrate, a Biodegradable Thermoplastic, Produced in
Transgenic Plants, "Science 256:520-523 (1992).
Pompon Denis et al., "Protein Engineering by cDNA Recombination in Yeasts: Shuffling of
Mamalian Cytochrome P-450 Functions, "Gene, 1989, vol. 83. pp. 15-24. Powell, K.T. et al., "Gel microdroplets and flow cytometry: Rapid determination of
antibody secretion by individual cells within a cell population, " Bio/Technology
8:333-337 (1990).
Rawlings, D.E. et al., "Mining with Microbes," Bio/Technology 13:773-778 (1995).
Rees, W.D. et al., "A molecular biological approach to reducing dietary amino acid
needs, "Bio/Technology 8:629-633 (1990).
Reidhaar-Olson et al., "Combinatorial Cassette Mutagenesis as a probe of the
Informational Content of Protein Sequences" Science 241:53-57 (1988).
Roa et al, Recombination and Polymerase Error Facilitate Restoration of Infectivity in
Brome Mosaic Virus Journal of Virology No. 2 67:969-979 (1993).
Sanseverino, J. et al., "Plasmid-Mediated Mineralization of Naphthalene, Phenanthrene,
Anthracene, "Applied Environ. Micro. 59(6):1931-1937 (1993).
Sarthy, A.V. et al., "Expression of the Escherichia coli Xylose Isomerase Gene in
Saccharomyces cerevisiae, "Appl. Environ. Micro. 53(9):1996-2000 (1987).
Scherrer, S. et al., "Periplasmic aggregation limits the proteolytic maturation of the
Escherichia coli Penicillin G amidase presursor polypeptide, "Appl. Microbiol.
Biotechnol. 42:85-91 (1994).
Selifonova, O.V. et al., "Use of an ibp-lux Fusion To Study Regulation of the
Isopropylbenzene Catabolism Operon of Pseudomonas putida RE204 and To Detect
Hydrophobic Pollutants in the Environment, "Appl. Environ. Microbiol. 62(3):778-783
(1986).
Silver, S. et al., "Newer Systems for Bacterial Resistances to Toxic Heavy Metals,"
Environ. Health Perspect. 102:107-113 (1994).
Simon, M.J. et al., "Sequences of genes encoding naphthalene dioxygenase in Pseudomonas
putida strains G7 and NCIB 9816-4, " Gene 127:31-37 (1993).
Simpson, T.W. et al., "Two paradigms of metabolic engineering applied to amino acid biosynthesis," Biochem. Soc. Trans. 23:381-387 (1995).
Smith, D.J. et al., "Cloning and heterologous expression of the penicillin biosynthetic
qene closuter from Penicillium chrysogenum, "Bio/Technology 8:39-41 (1990).
Steele, D.B. et al., "Techniques for selection of industrially important
microorganisms, " Ann. Rev. Microbiol. 45:89-106 (1991).
Stemmer, W.P.C., "Rapid Evolution of a Protein In Vitro by DNA Shuffling," Nature
370:389-391 (1994).
Stemmer, W.P.C., "DNA Shuffling by Random Fragmentation and Reassembly: In Vitro
Recombination for Molecular Evolution, " Proc. Natl. Acad. Sci. U.S.A. 91:10747-10751
(1994).
Stemmer et al., "Selection of an Active Single Chain FV Antibody From a Protein Linker
Library Prepared by Enzymatic Inverse PCR" Biotechniques 14:256-265 (1992).
Stephanopoulos, G. et al., "Metabolic engineering--methodologies and future prospects,"
Trends. Biotech. 11:392-396 (1993).
Stephanopoulos, G., "Metabolic engineering," Curr. Opin. Biotech. 5:196-200 (1994).
Tabata, K. et al., "A carotenogenic gene cluster exists on a large plasmid in Thermus
thermophilus, "FEBS Letts. 341:251-255 (1994).
Timmis, K.N. et al., "Designing microorganisms for the Treatment of Toxic Wastes," Ann.
Rev. Microbiol. 48:525-557 (1994).
Wackett, L.P. et al., "Metabolism of polyhalogenated compounds by a genetically engineered bacterium," Nature 368:627-629 (1994).
Wang et al., "Identification of Glial Filament Protein and Vimentin in the Same
Intermediate Filament System in Human Glioma Cells, Proc. Natl. Sci. USA
81(7):2102-2106 (Abstract only, 1984).
Wang, Y et al., "Nucleotide Sequence of a Chromosomal Mercury Resistance Determinant
from a Bacillus sp. With Broad-Specrum Mercury Resistance, " J. Bacteriol. 171(1):83-92
Weber, H. et al., "Formation of Genes Coding for Hybrid Proteins by Recombination
Between Related, Cloned Genes in E. coli, "Nucleic Acids Research 11(16): 5661-5669
Weaver, J.C. et al., "Microdrop Technology: A General Method for Separating Cells by
Function and Composition, "Methods 2:234-247 (1991).
Wehmeier, U.F., "New multifunctional Escherichia coli-Streptomyces shuttle vectors
allowing blue-white screening on Xgal plates, "Gene 165:149-150 (1995).
```

```
Zhao, X., "EPD, a novel technology for drug delivery," Adv. Drug Delivery Rev.
17:257-262 (1995).
Zukowski, M.M. et al., "Chromogenic identification of genetic regulatory signals in
Bacillus subtilis based on expression of a cloned Pseudomonas gene, " Proc. Natl. Acad.
Sci. U.S.A. 80:1101-1105 (1983).
Zylstra, G.J. et al., "Cloning and Analysis of the Genes for Polycyclic Aromatic
Hydrocarbon Degradation, "Ann. NY Acad. Sci. 721:386-398 (1994).
Crameri a et al: "Molecular evolution of an arsenate detoxification pathway by DNA
shuffling" Nature Biotechnology, vol.15, May 1997, pp. 436-438, XP002082183.
Clackson et al., "Making antibody fragments using phage display libraries," Nature,
352:624-628 (Aug. 15, 1991).
Crameri et al., "10(20)-Fold aptamer library amplification without gel purification," Nuc. Acids Res., 21(18):4410 (1993).
Cull et al., "Screening for receptor ligands using large libraries of peptides linked
to the C terminus of the lac repressor, PNAS, 89:1865-1869 (Mar. 1992).
Cwirla et al., "Peptides on phage: A vast library of peptides for identifying ligands,".
PNAS, 87:6378-6382 (Aug. 1990).
Daugherty et al., "Polymerase chain reaction facilitates the cloning, CDR-grafting, and
rapid expression of a murine monoclonal antibody directed against the CD18 component of
leukocyte integrins, " Nuc. Acids Res., 19(9):2471-2476 (1991).
Delagrave et al., "Searching Sequence Space to Engineer Proteins: Exponential Ensemble
Mutagenesis, Biotechnology, 11:1548-1552 (Dec. 1993).
Dube et al., "Artificial mutants Generated by the Insertion of Random Oligonucleotides
into the Putative Nucleoside Binding Site of the HSV-1 Thymidine Kinase Gene,"
Biochemistry, 30(51):11760-11767 (1991).
Ghosh et al., "Arginine-395 Is Required for Efficient in Vivo and in Vitro
Aminoacylation of tRNAs by Escherichia coli Methionyl-tRNA Stnthetase, "Biochemistry,
30:11767-11774 (1991).
Goldman et al., "An Algorithmically Optimized Combinatorial Library Screened by digital
Imaging Spectroscopy, "Biotechnology, 10:1557-1561 (Dec. 1992).
Harlow et al., "Construction of Linker-Scanning Mutations using the Polymerase Chain
Reaction, " Methods in Mol. Biol., 31:87-96 (1994).
Heda et al., "A simple in vitro site directed mutagenesis of concatamerized cDNA by
inverse polymerase chain reaction, " Nuc. Acids Res., 20(19):5241-5242 (1992)
Ho et al., "DNA and Protein Engineering Using the Polymerase Chain Reaction: Splicing
by Overlap Extension, DNA and Protein Eng. Techniques , 2(2):50-55 (1990).
Hodgson, "The Whys and Wherefores of DNA Amplification," Biotechnology, 11:940-942
(Aug. 1993).
Horton et al., "Gene Splicing by Overlap Extension," Mehtods in Enzymology, 217:270-279
(1993).
Horton et al., "Gene Splicing by Overlap Extension: Tailor-Made Genes Using the
Polymerase chain Reaction, "BioTechniques, 8(5):528-535 (May 1990).
Jayaraman et al., "Polymerase chain reaction-mediated gene synthesis: Synthesis of a
gene coding for isozyme c of horseradish peroxidase, PNAS, 88:4084-4088 (May 1991).
Jones et al., "A Rapid Method for Recombination and Site-Specific Mutagenesis by
Placing Homologous ends on DNA Using Polymerase Chain Reaction, "BioTechniques, 10(1):
62-66 (1991).
Joyce, G. F., "Directed Molecular Evolution," Scientific American , (Dec. 1992). Klug et al., "Creating chimeric molecules by PCR directed homologous DNA
recombination, " Nuc. Acids Res., 19(10):2793 (1991).
Krishnan et al., "Direct and crossover PCR amplification to facilitate Tn5supF-based
sequencing of .lambda. phage clones, " Nuc. Acids Res., 19(22):6177-6182 (1991).
Majumder, K., "Ligation-free gene synthesis by PCR: synthesis and mutagenesis at
multiple loci of a chimeric gene encoding OmpA signal peptide and hirudin, " Gene,
110:89-94 (1992).
Marks et al., "By-passing Immunization, Human Antibodies from V-gene Libraries
Displayed on Phage, " J. Mol. Biol., 222:581-597 (1991).
McCafferty et al., "Phage antibodies: filamentous phage displaying antibody variable
domains, " Nature, 348:552-554 (Dec. 6, 1990).
Morl et al., "Group II intron RNA-catalyzed recombination of RNA in vitro," Nuc. Acids
Res., 18(22):6545-6551 (1990).
Mullis et al., "Specific Synthesis of DNA in Vitro via a Polymerase-Catalyzed Chain
Reaction, "Methods in Enzymology, 155:335-351 (1987).
Mullis et al., "Specific Enzymatic Amplification of DNA In Vitro: The Polymerase Chain
Reaction, "Cold Spring Harbor Symposia on Quantitative Biology, 51:263-273 (1986). Nissim et al., "Antibody fragments from a `single pot` display library as
immunochemical reagents," EMBO Journal, 13(3):692-698 (1994).
Osuna et al., "Combinatorial mutagenesis of three major groove-contacting residues of
Eco RI: single and double amino acid replacements retaining methyltransferase-sensitive
```

```
activities, " Gene, 106:7-12 (1991).
Paabo et al., "DNA Damage Promotes Jumping between Templates during Enzymatic
Amplification, J. Biol. Chem., 265(8):4718-4721 (Mar. 15, 1990).
Saiki et al., "Diagnosis of sickle Cell Anemia and .beta.-Thalassemia with
Enzymatically Amplified DNA and Nonradioactive Allele-Specific Oligonucleotide Probes,"
New England J. of Medicine, 319(9):537-541 (Sep. 1, 1988).
Saiki et al., "analysis of enzymatically amplified .beta.-globin and HLA-DQ.alpha. DNA
with allele-specific oligonucleotide probes, " Nature, 324:163-166 (Nov. 13, 1986).
Saiki et al., "Enzymatic Amplification of .beta.-Globin Genomic Sequences and Restriction Site analysis for Diagnosis of Sickle Cell Anemia," Science, 230:1350-1354
(Dec. 20, 1985).
Saiki et al., "Primer-Directed Enzymatic Amplification of DNA with a Thermostabl; e DNA
Polymerase, "Science, 239:487-491 (Jan. 20, 1988).
Sambrook et al., Molecular Cloning, A Laboratory Manual, Cold Spring Laboratory Press,
Cold Spring Harbor, New York (1989).
Scharf et al., "Direct Cloning and Sequence Analysis of Enzymatically Amplified Genomic
Sequences, " Science, 233:1076-1078 (Sep. 1986).
Scott et al., "Searching for Peptide Ligands with an Epitope Library," Science,
249:386-390 (Jul. 20, 1990).
Sikorski et al., "In Vitro Mutagenesis and Planned Shuffling: From Cloned Gene to
Mutant Yeast, "Methods in Enzymology, 194:302-318 (1991).
Smith et al., "Unwanted Mutations in PCR Mutagenesis: Avoiding the Predictable," PCR
Methods and Applications, 2(3):253-257 (Feb. 1993).
Villarreal et al., "A General Method of Polymerase-Chain-Reaction-Enabled Protein
Domain Mutagenesis: Construction of a Human Protein S-Osteonectin Gene, " Analytical
Biochem., 197:362-367 (1991).
Weisberg et al., "Simultaneous Mutagenesis of Multiple Sites: Application of the Ligase
Chain Reaction Using PCR Products Instead of Oligonucleotides, "BioTechnques,
15(1):68-70, 72-74, 76 (Jul. 1993).
Weissenhorn et al., "Chimerization of antibodies by isolation of rearranged genomic
variable regions by the polymerase chain reaction, " Gene, 106:273-277 (1991).
Yao et al., "Site-directed Mutagenesis of Herpesvirus Glycoprotein Phoshphorylation
Sites by Recombination Polymerase Chain Reaction, " PCR Methods and Applications,
1(3):205-207 (Feb. 1992).
Yolov et al., "Constructing DNA by polymerase recombination," Nuc. Acids Res.,
18(13):3983-3986 (1990).
Yon et al., "Precise gene fusion by PCR," Nuc. Acids Res., 17(12):4895 (1989).
Zoller, M.J., "New recombinant DNA methodology for protein engineering," Curr. Opin.
Biotech., 3:348-354 (1992).
Opposition Statement in matter of Australian Patent Application 703264 (Affymax
Technologies NV), filed by Diversa Corporation on Sep. 23, 1999.
Andersson et al., "Muller's ratchet decreases fitness of a DNA-based microbe", PNAS, 93: 906-907 (Jan. 1996).
Bailey, "Toward a Science of Metabolic Engineering", Science, 252: 1668-1680 (1991).
Barrett et al., "Genotypic analysis of multiple loci in somatic cells by whole genome
amplification", Nuc. Acids Res., 23(17): 3488-3492 (1995).
Cameron et al., "Cellular and Metabolic Engineering An Overview", Applied Biochem.
Biotech., 38: 105-140 (1993).
Chakrabarty, "Microbial Degradation of Toxic Chemicals: Evolutionary Insights and
Practical Considerations", ASM News, 62(3): 130-137 (1996).
Chater, "The Improving Prospects for Yield Increase by Genetic Engineering in
Antibiotic-Producing Streptomycetes", Biotechnology, 8: 115-121 (Feb. 1990).
Chen et al., "Tuning the activity of an enzyme for unusual environments: Sequential
random mutagenesis of subtilisin E for catalysis in dimethylformamide", PNAS, 90:
5618-5622 (Jun. 1993).
Dieffenbach et al., PCR Primer, A Laboratory Manual, Cold Spring Harbor Laboratory Press, pp. 583-589, 591-601, 603-612, and 613-621 (1995).
Evnin et al., "Substrate specificity of trypsin investigated by using a genetic
selection", PNAS, 87: 6659-6663 (Sep. 1990).
Fang et al., "Human Strand-specific Mismatch Repair Occurs by a Bidirectional Mechanism
Similar to That of the Bacterial Reaction", J. Biol. Chem., 268(16): 11838-11844 (Jun.
5, 1993).
Ippolito et al., "Structure assisted redesign of a protein-zinc-binding site with
femtomolar affinity", PNAS, 92: 5017-5021 (May 1995).
Kellogg et al., "Plasmid-Assisted Molecular Breeding: New Technique for Enhanced
Biodegradation of Persistent Toxic Chemicals", Science, 214: 1133-1135 (Dec. 4, 1981).
Kunkel, "Rapid and efficient site-specific mutagenesis without phenotypic selection",
PNAS, 82: 488-493 (Jan. 1985).
Levichkin et al., "A New Approach to Construction of Hybrid Genes: Homolog
```

```
Recombination Method", Mol. Biology, 29(5) part 1: 572-577 (1995).
Lewis et al., "Efficient site directed in vitro mutagenesis using ampicillin
selection", Nuc. Acids Res., 18(12): 3439-3443 (1990).
Moore et al., "Directed evolution of a para-nitrobenzyl esterase for aqueous-organic
solvents", Nature Biotech., 14: 458-467 (Apr. 1996).
Omura, "Philosophy of New Drug Discovery", Microbiol. Rev., 50(3): 259-279 (Sep. 1986).
Piepersberg, "Pathway Engineering in Secondary Metabolite-Producing Actinomycetes",
Crit. Rev. Biotech., 14(3):251-285 (1994).
Prasher, "Using GFP to see the light", TIG, 11(8): (Aug. 1995).
Rice et al., "Random PCR mutagenesis screening of secreted proteins by direct
expression in mammalian cells", PNAS, 89: 5467-5471 (Jun. 1992).
Simpson et al., "Two paradigms of metbolic engineering applied to amino acid
biosynthesis", Biochem. Soc. Transactions, vol. 23 (1995).
Steele et al., "Techniques for Selection of Industrially Important Microorganisms",
Ann. Rev. Microbiol., 45: 89-106 (1991). Stephanopoulos et al., "Metabolic engineering--methodologies and future prospects",
Trends Biotech. 11: 392-396 (1993).
Stephanopoulos, "Metabolic engineering", Curr. Opin. Biotech., 5: 196-200 (1994).
Wehmeier, "New multifunctional Escherichia coli-Streptomyces shuttle vectors allowing
blue-white screening on XGal plates", Gene, 165: 149-150 (1995).
Fisch et al., "A Strategy Of Exon Shuffling For Making Large Peptide Repertoires
Displayed On Filamentous Bacteriophage", Proc Natl Acad Sci USA, 93(15):7761-7766
(1996).
Marton et al., "DNA Nicking Favors PCR Recombination", Nucleic Acids Res.,
19(9):2423-2426 (1991).
Winter et al., "Making Antibodies By Phage Display Technology", Ann. Rev. Immunol.,
12:433-455 (1994).
Greener et al., "An Efficient Random Mutagenesis Technique Using An E. coli Mutator
Strain", Methods in Molecular Biology, 57:375-385 (1995).
Balint et al., "Antibody Engineering By Parsimonious Mutagenesis", Gene, 137(1):109-118
(1993).
Bartel et al., "Isolation of New Ribozymes From a Large Pool of Ramdon Sequences",
Science, 261:1411-1418 (1993).
Crameri et al., "Combinatorial Multiple Cassette Mutagenesis Creates All The
Permutations Of Mutant And Wild-Type Sequences", Biotechniques, 18(2):194-196 (1995).
Crameri et al., "Improved Green Fluorescent Protein By Molecular Evolution Using DNA
Shuffling", Nat. Biotechnol., 14(3):315-319 (1996).
Crameri et al., "Construction And Evolution Of Antibody-Phage Libraries By DNA
Shuffling", Nat. Med., 2(1):100-102 (1996).
Crameri et al., "Molecular Evolution Of An Arsenate Detoxification Pathway By DNA Shuffling", Nat. Biotechnol., 15(5):436-438 (1997).
Crameri et al., "DNA Shuffling Of A Family Of Genes From Diverse Species Accelerates
Directed Evolution", Nature, 391(3664):288-291 (1998).
Gates et al., "Affinity Selective Isolation Of Ligands From Peptide Libraries Through
Display On A Lac Repressor `Headpiece Dimer`", J. Mol. Biol., 255(3):373:386 (1996).
Gram et al., "In Vitro Selection and Affinity Maturation of Antibodies From a Naive
Combinatorial Immunoglobulin Library", Proc. Natl. Acad. Sci. USA, 89:3576-3580 (1992).
Near, "Gene Conversion Of Immunoglobulin Variable Regions In Mutagenesis Cassettes By
Replacement PCR Mutagenesis", Biotechniques. 12(1):88-97 (1992).
Perlak, "Single Step Large Scale Site-Directed In Vitro Mutagenesis Using Multiple
Oligonucleotides", Nucleic Acids Res., 18(24):7457-7458 (1990).
Stemmer, "Searching Sequence Space", Biotechnology. 13:549-553 (1995).
Stemmer et al., "Single-Step Assembly Of A Gene And Entire Plasmid From Large Numbers
Of Oligodeoxyribonucleotides". Gene, 164(1):49-53 (1995).

Stemmer, "The Evolution of Molecular Computation". Science, 270(5241):1510 (1995).

Stemmer, "Sexual PCR and Assembly PCR" Encyclopedia Mol. Biol., VCH Publishers, New
York, pp. 447-457 (1996).
Weber et al., "Formation of Genes Coding for Hybrid Proteins by Recombination Between
Related, Cloned Genes in E. coli, "Nucleic Acids Research, 11(16):5661-5669 (1983).
Weisberg et al., "Simultaneous Mutagenesis Of Multiple Sites: Application Of The Ligase
Chain Reaction Using PCR Products Instead Of Oligonucleotides". BioTechniques,
15(1):68-76 (1993).
Zhang et al., "Directed Evolution Of A Fucosidase From A Galactosidase By DNA Shuffling
And Screening", Proc. Natl. Acad. Sci. USA, 94(9):4504-4509 (1997).
Zhao et al., "Molecular Evolution by Staggered Extension Process (StEP) In Vitro
Recombination", Nature Biotech., 16:258-261 (1998).
Arkin et al., "An algorithm for protein engineering: Stimulation of recursive ensemble
```

1/14/03 1:16 PM

```
mutagenesis" Proc. Natl. Acad. Sci. USA 89:7811-7815 (1992).
Beaudry et al., "Directed evolution of an RNA enzyme" Science 257:635-641 (1992).
Berger et al. "Phoenix mutagenesis: One-step reassembly of multiply cleaved plasmids
with mixtures of mutant and wild-type fragments" Anal. Biochem. 214:571-579 (1993).
Berkhout et al. "In vivo selection of randomly mutated retroviral genomes" Nucleic
Acids Res. 21:5020-5023 (1993).
Cadwell et al. "Randomization of genes by PCR mutagenesis" PCR Methods and Applications
2:28-33 (1992).
Calogero et al. "In vivo recombination and the production of hybrid genes" Microbiol.
Lett. 97:41-44 (1992).
Caren et al. "Efficient sampling of protein sequence space for multiple mutants"
Biotechnology 12:517-520 (1994).
Delagrave et al. "Recursive ensemble mutagenesis" Protein Engineering 6:327-331 (1993).
Feinberg and Vogelstein "A technique for radiolabeling DNA restriction endonuclease
fragments to high specific activity" Anal. Biochem. 132:6-13 (1983).
Heim et al. "Wavelength mutations and postranslational autoxidation of green
fluorescent protein Proc. Natl. Acad. Sci. USA 91:12501-12504 (1994).
Hermes et al. "Searching sequence space by definably random mutagenesis: Improving the
catalytic potency of an enzyme" Proc. Natl. Acad. Sci. USA 87:696-700 (1990).
Ho et al. "Site-directed mutagenesis by overlap extension using the polymerase chain
reaction" Gene 77:51-59 (1989).
Horton et al. "Engineering hybrid geneswithout the use of restriction enzymes: Gene
splicing by overlap extension" Gene 77:61-68 (1989).
Jones et al. "Recombinant cicle PCR and recombination PCR for site-specific mutagenesis
without PCR product purification" BioTechniques 12:528-530, 532, 534-535 (1992).
Kim et al. "Human immunodeficiency virus reverse transcriptase" J. Biol. Chem.
271:4872-4878 (1996).
Leung et al. "A method for random mutgenesis of a defined DNA segment using a modified
polymerase chain reaction" Techniques 1:11-15 (1989).
Marks et al. "By-passing immunization: Building high affinity human antibosies by chain
shuffling" Bio/Technology 10:779-782 (1992).
Meyerhans et al. "DNA recombination using PCR" Nucleic Acids Res. 18:1687-1691 (1990).
Oliphant et al. "Cloning of random-sequence oligodeoxynucleotides" Gene 44:177-183
(1988).
Pharmacia Catalog pp. 70-71 (1993 Edition).
Pompon et al. "Protein engineering by cDNA recombination in yeasts: Shuffling of
mammalian cytochrome P-450 functions" Gene 83:15-24 (1989).
Reidhaar-Olson et al. "Combinatorial cassette mutagenesis as a probe of the
informational content of protein sequences" Science 241:53-57 (1988).
Roa et al. "Recombination and polymerase error facilitate restoration of infectivity in
brome mosaic virus" J. Virol. 67:969-979 (1993).
Stemmer et al. "Selection of an active single chain Fv antibody from a protein linker
library prepared by enzymatic inverse PCR" BioTechniques 14:256-265 (1992).
Stemmer et al. "Rapid evolution of a protein in vitro by DNA shuffling" Nature
370:389-391 (1994).
Stemmer et al. "DNA shuffling by random fragmentation and reassembly: In vitro
recombination for molecular evolution" Proc. Natl. Acad. Sci. USA 91:10747-10751
(1994).
Dieffenbach et al., PCR Primer, A Laboratory Manual, New York, Cold Spring Harbor Press, pp. 581-621 (1995).
Higuchi, R., "Using PCT to Engineer DNA," PCR Technology, Erlich, H.A., ed., New York,
Stockton Press, pp. 61-70 (1989).
Andersson, D. et al., "Muller's Ratchet Decreases Fitness of a DNA-based Microbe," Proc. Natl. Acad. Sci. U.S.A. 93:906-907 (1996).
Barrett, M. et al., "Genotypic Analysis of Multiple Loci in Somatic Cells by Whole
Genome Amplification, "Nucleic Acids Research 23:3488-3492 (1995).
Crameri, A. et al., "Improved Green Fluorescent Protein by Molecular Evolution Using
DNA Shuffling," Nature Biotechnology 14:315-319 (1996).
Crameri, A. et al., "Construction and Evolution of Antibody-phage Libraries by DNA
Shuffling, "Nature Medicine 2:100-300 (1996).
Feinberg, A. et al., "A Technique for Radiolabeling DNA Restriction Endonuclease
Fragments to High Specific Activity, " Analytical Biochemistry 132:6-13 (1983).
Ho, S. et al., "Site-directed Mutagenesis by Overlap Extension Using the Polymerase
Chain Reaction, Gene 77:51-59 (1989).
Horton, R. et al., "Engineering Hybrid Genes Without the Use of Restriction Enzymes:
Gene Splicing by Overlap Extension, " Gene 77:61-68 (1989).
Levichkin, I.V. et al., "A New Approach to Construction of Hybrid Genes: Homolog
Recombination Method, " Molecular Biology 29:572-577 (1995).
```

Pharmacia Catalog, "Recombinant Phage Antibody System: Mouse ScFv Module Expression Module Detection Module, p. 70-71, 1993 Edition.
Ramos, J.L. et al., "The Behavior of Bacteria Designed for Biodegradation,"

Bio/Technology 12:1349-1355 (1994). Stemmer, W.P.C. et al., "Single-step Assembly of a Gene and Entire Plasmid from Large Numbers of Oligodeoxyribonucleotides, "Gene 164:49-53 (1995).

Stemmer, W.P.C., "Searching Sequence Space: Using Recombination to Search More Efficiently and Thoroughly Instead of Making Bigger Combinatorial Libraries," Bio/Technology 13:549-553 (1995).

Stemmer, W.P.C., "Rapid Evolution of a Protein In Vitro by DNA Shuffling," Nature 370:389-391 (1994).

Stemmer, W.P.C., "The Evolution of Molecular Computation," Science 270:1510 (1995). Stemmer, W.P.C., "DNA Shuffling by Random Fragmentation and Reassembly: In Vitro Recombination for Molecular Evolution, Proc. Natl. Acad. Sci. U.S.A. 91:10747-10751 (1994).

ART-UNIT: 165

PRIMARY-EXAMINER: Whisenant; Ethan

ABSTRACT:

The present invention is generally directed to the evolution of new metabolic pathways and the enhancement of bioprocessing through a process herein termed recursive sequence recombination. Recursive sequence recombination entails performing iterative cycles of recombination and screening or selection to "evolve" individual genes, whole plasmids or viruses, multigene clusters, or even whole genomes. Such techniques do not require the extensive analysis and computation required by conventional methods for metabolic engineering.

134 Claims, 1 Drawing figures

Generate Collection

Print

Search Results - Record(s) 1 through 10 of 20 returned.

1. Document ID: US 6488850 B2

L2: Entry 1 of 20

File: USPT

Dec 3, 2002

US-PAT-NO: 6488850

DOCUMENT-IDENTIFIER: US 6488850 B2

TITLE: Method and apparatus for anaerobically degrading pollutants with alkanes

DATE-ISSUED: December 3, 2002

INVENTOR-INFORMATION:

NAME

CITY

ZIP CODE STATE

COUNTRY

Perriello; Felix Anthony

Norwood MA

US-CL-CURRENT: 210/605; 210/170, 210/220, 210/611, 210/747, 210/908, 435/262, 435/262.5

Full Title Citation Front Review Classification Date Reference Sequences Attachments Claims Draw, Desc Image

2. Document ID: US 6245235 B1

L2: Entry 2 of 20

File: USPT

Jun 12, 2001

US-PAT-NO: 6245235

DOCUMENT-IDENTIFIER: US 6245235 B1

TITLE: System and method of in-situ bioremediation with butane-utilizing bacteria

DATE-ISSUED: June 12, 2001

INVENTOR-INFORMATION:

NAME

CITY

STATE

ZIP CODE

COUNTRY

Perriello; Felix Anthony

Norwood

MA

02062

US-CL-CURRENT: 210/611; 210/620, 210/747, 210/908, 210/909, 435/262.5

Title Citation Front Review Classification Date Reference Sequences Attachments Claims KWC Draw, Desc Image

3. Document ID: US 6241779 B1

L2: Entry 3 of 20

File: USPT

Jun 5, 2001

US-PAT-NO: 6241779

ZIP CODE

Record List Display

DOCUMENT-IDENTIFIER: US 6241779 B1

TITLE: Metal ligand containing bleaching compositions

DATE-ISSUED: June 5, 2001

INVENTOR-INFORMATION:

NAME CITY STATE

Collins; Terrence J. Pittsburgh PA Horwitz; Colin P. Pittsburgh PA

 $\text{US-CL-CURRENT: } \underline{8/111}; \ \underline{252/186.33}, \ \underline{252/186.39}, \ \underline{252/186.43}, \ \underline{510/311}, \ \underline{8/107}, \ \underline{8/108.1}$

Full | Title | Citation | Front | Review | Classification | Date | Reference | Sequences | Attachments |
Draw, Desc | Image |

KMC

COUNTRY

4. Document ID: US 6210579 B1

L2: Entry 4 of 20 File: USPT

Apr 3, 2001

US-PAT-NO: 6210579

DOCUMENT-IDENTIFIER: US 6210579 B1

TITLE: Bioremediation of pollutants with butane-utilizing bacteria

DATE-ISSUED: April 3, 2001

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Perriello; Felix Anthony West Roxbury MA

US-CL-CURRENT: 210/611; 210/620, 210/747, 210/908, 210/909, 435/262.5

Full | Title | Citation | Front | Review | Classification | Date | Reference | Sequences | Attachments |
Draw, Desc | Image |

KWIC

5. Document ID: US 6136223 A

L2: Entry 5 of 20

File: USPT Oct 24, 2000

US-PAT-NO: 6136223

DOCUMENT-IDENTIFIER: US 6136223 A

TITLE: Metal ligand containing bleaching compositions

DATE-ISSUED: October 24, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Collins; Terrence J. Pittsburgh PA Horwitz; Colin P. Pittsburgh PA

US-CL-CURRENT: 252/186.33; 252/186.39, 252/186.43, 510/311, 540/460, 540/465

Full | Title | Citation | Front | Review | Classification | Date | Reference | Sequences | Attachments | Draw, Desc | Image |

KWIC

6. Document ID: US 6110372 A

L2: Entry 6 of 20

File: USPT

Aug 29, 2000

US-PAT-NO: 6110372

DOCUMENT-IDENTIFIER: US 6110372 A

TITLE: Bioremediation of petroleum pollutants with alkane-utilizing bacteria

DATE-ISSUED: August 29, 2000

INVENTOR-INFORMATION:

NAME

CITY

STATE

ZIP CODE

COUNTRY

Perriello; Felix Anthony

Norwood

AM

02062

ZIP CODE

US-CL-CURRENT: 210/611; 210/620, 210/747, 210/908, 210/909, 435/262, 435/262.5

Full Title Citation Front Review Classification Date Reference Sequences Attachments

Draws Description

KWIC

7. Document ID: US 6107528 A

L2: Entry 7 of 20

File: USPT

Aug 22, 2000

US-PAT-NO: 6107528

DOCUMENT-IDENTIFIER: US 6107528 A

TITLE: Iron complexes for bleach activation and stereospecific oxidation

DATE-ISSUED: August 22, 2000

INVENTOR-INFORMATION:

NAME

CITY

STATE

COUNTRY

Que, Jr.; Lawrence

Roseville

MN

Kim; Cheal

Minneapolis

MN

rem, chear

Zanq; Yan

Chapel Hill

Kim; Jinheung

Minneapolis

NC MN

US-CL-CURRENT: $\underline{568}/\underline{832}$; $\underline{252}/\underline{186.38}$, $\underline{252}/\underline{186.39}$, $\underline{252}/\underline{186.4}$, $\underline{252}/\underline{186.41}$, $\underline{252}/\underline{186.42}$, $\underline{510}/\underline{311}$, $\underline{510}/\underline{376}$, $\underline{556}/\underline{138}$, $\underline{8}/\underline{111}$

Full Title Citation Front Review Classification Date Reference Sequences Attachments

Draw Desc Image

KWIC

8. Document ID: US 6100394 A

L2: Entry 8 of 20

File: USPT

Aug 8, 2000

US-PAT-NO: 6100394

DOCUMENT-IDENTIFIER: US 6100394 A

TITLE: Long-lived homogenous oxidation catalysts

DATE-ISSUED: August 8, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Collins; Terrence J. Pittsburgh PA Gordon-Wylie; Scott W. Pittsburgh PA

US-CL-CURRENT: 540/467; 502/150, 540/450, 540/451, 540/452, 540/453, 540/465, 540/480,

<u>540/482</u>, <u>540/483</u>

Full | Title | Citation | Front | Review | Classification | Date | Reference | Sequences | Attachments |
Draw, Desc | Image |

9. Document ID: US 6099586 A

L2: Entry 9 of 20 File: USPT Aug 8, 2000

US-PAT-NO: 6099586

DOCUMENT-IDENTIFIER: US 6099586 A

TITLE: Metal ligand containing bleaching compositions

DATE-ISSUED: August 8, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Collins; Terrence J. Pittsburgh PA Horwitz; Colin P. Pittsburgh PA

US-CL-CURRENT: 8/111; 252/186.33, 252/186.39, 252/186.43, 510/311, 8/108.1

Full Title Citation Front Review Classification Date Reference Sequences Attachments KMC |

10. Document ID: US 6054580 A

L2: Entry 10 of 20 File: USPT Apr 25, 2000

US-PAT-NO: 6054580

DOCUMENT-IDENTIFIER: US 6054580 A

TITLE: Long-lived homogenous amide containing macrocyclic compounds


DATE-ISSUED: April 25, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Collins; Terrence J. Pittsburgh PA
Gordon-Wylie; Scott W. Burlington VT
Horwitz; Colin P. Pittsburgh PA

US-CL-CURRENT: 540/460; 540/465, 540/480, 540/482, 540/483



Display Format: CIT Change Format

Previous Page Next Page

WEST

Generate Collection

Print

Search Results - Record(s) 1 through 2 of 2 returned.

1. Document ID: US 6117661 A

L9: Entry 1 of 2

File: USPT

Sep 12, 2000

US-PAT-NO: 6117661

DOCUMENT-IDENTIFIER: US 6117661 A

TITLE: Mutant mono-oxygenase cytochrome P450cam

DATE-ISSUED: September 12, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Wong; Luet-Lok Oxford GB
Flitsch; Sabine Lahja Edinburgh GB
Nickerson; Darren Paul Oxford GB
Hart; Alwyn James Loughborough GB

US-CL-CURRENT: 435/189; 435/252.3, 435/320.1, 435/471, 435/69.1, 536/23.2

Full Title Citation Front Review Classification Date Reference Sequences Attachments

Draw. Desc. | Image |

KWIC

2. Document ID: US 6100074 A

L9: Entry 2 of 2

File: USPT

Aug 8, 2000

US-PAT-NO: 6100074

DOCUMENT-IDENTIFIER: US 6100074 A

TITLE: Mutant mono-oxygenase cytochrome P-450 .sub.cam

DATE-ISSUED: August 8, 2000

INVENTOR - INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Flitsch; Sabine Lahja Edinburgh GB
Nickerson; Darren Paul Oxford GB
Wong; Luet-Lok Oxford GB

US-CL-CURRENT: 435/189; 435/132, 435/69.1, 530/402

Full Title Citation Front Review Classification Date Reference Sequences Attachments
Draw, Desc Image


KWIC

Generate Collection Print

Terms	Documents
mutant mono-oxygenase	2

Display Format: CIT Change Format

<u>Previous Page</u> <u>Next Page</u>



L9: Entry 1 of 2

File: USPT

Sep 12, 2000

US-PAT-NO: 6117661

DOCUMENT-IDENTIFIER: US 6117661 A

TITLE: Mutant mono-oxygenase cytochrome P450cam

DATE-ISSUED: September 12, 2000

INVENTOR-INFORMATION:

CITY	STATE	ZIP	CODE	COUNTRY
Oxford				GB
Edinburgh				GB
Oxford				GB
Loughborough				GB
	Oxford Edinburgh Oxford	Oxford Edinburgh Oxford	Oxford Edinburgh Oxford	Oxford Edinburgh Oxford

US-CL-CURRENT: 435/189; 435/252.3, 435/320.1, 435/471, 435/69.1, 536/23.2

CLAIMS:

What is claimed is:

- 1. A <u>mutant mono-oxygenase</u> cytochrome P450cam comprising either a deletion of the cysteine at amino acid position 334 or a substitution of another amino acid for the cysteine at amino acid position 334.
- 2. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 1 wherein an amino acid other than cysteine is substituted at amino acid position 334.
- 3. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 2 further comprising the substitution of an amino acid other than tyrosine at amino acid position 96.
- 4. A <u>mutant mono-oxygenase</u> cytochrome P.sup.450 cam according to claim 2 further comprising one or more amino acid substitutions at amino acid positions selected from the group consisting of 87, 98, 101, 185, 193, 244, 247, 295, 297, 395, and 396.
- 5. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 1 wherein the cysteine is deleted at amino acid position 334.
- 6. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 5 further comprising the substitution of an amino acid other than tyrosine at amino acid position 96.
- 7. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 5 further comprising one or more amino acid substitutions at amino acid positions selected from the group consisting of 87, 98, 101, 185, 193, 244., 247,

295, 297, 395, and 396.

8. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 1 further comprising the substitution of an amino acid other than tyrosine at amino acid

position 96.

- 9. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 8 wherein the substituent amino acid for position 96 is selected from the group consisting of alanine, arginine, asparagine, aspartic acid, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, proline, serine, threonine, tryptophan, tyrosine and valine.
- 10. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 8 further comprising one or more amino acid substitutions at amino acid positions selected from the group consisting of 87, 98, 101, 185, 193, 244, 247, 295, 297, 395, and 396.
- 11. A mutant mono-oxgenase cytochrome P450cam according to claim 1 wherein the substituent amino acid is selected from the group consisting of alanine, arginine, asparagine, aspartic acid, glutamic acid, glutamine, glycine, histidine, isoleucilne, leucine, lysine, methionine, proline, serine, threonine, tryptophan, tyrosine and valine.
- 12. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 11 further comprising one or more amino acid substitutions at amino acid positions selected from the group consisting of 87, 98, 101, 185, 193, 244, 247, 295, 297, 395, and 396.
- 13. A <u>mutant mono-oxygenase</u> cytochrome P450cam according to claim 1 further comprising one or more amino acid substitutions at amino acid positions selected from the group consisting of 87, 98, 101, 185, 193, 244, 247, 295, 297, 395, and 396.

WEST	

End of Result Set

	\$0000000000000000000000000000000000000	3 commence	٤
	Generate Collection	Print	
3039990	Ł	£	3

L9: Entry 2 of 2

File: USPT

Aug 8, 2000

US-PAT-NO: 6100074

DOCUMENT-IDENTIFIER: US 6100074 A

TITLE: Mutant mono-oxygenase cytochrome P-450 .sub.cam

DATE-ISSUED: August 8, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY Flitsch; Sabine Lahja Edinburgh GB

Nickerson; Darren Paul Oxford GB
Wong; Luet-Lok Oxford GB

US-CL-CURRENT: 435/189; 435/132, 435/69.1, 530/402

CLAIMS:

We claim:

- 1. A <u>mutant mono-oxygenase</u> cytochrome P-450.sub.cam wherein the tyrosine residue at position 96 is replaced by the residue of a small hydrophobic amino acid.
- 2. The mutant of claim 1, wherein said mutant catalyzes the oxidation of a compound selected from the group consisting of a polycyclic aromatic hydrocarbon, a linear or branched alkane, a biphenyl compound and a halogenated hydrocarbon.
- 3. The mutant of claim 1, wherein the amino acid is selected from the group consisting of alanine, glycine, isoleucine, leucine, and valine.
- 4. The mutant of claim 1, wherein an amino acid residue at one or more of the positions 87, 98, 185, 244, 247, 295 or 297 is independently replaced by another amino acid residue.
- 5. The mutant of claim 2, wherein the amino acid is selected from the group consisting of alanine, glycine, isoleucine, leucine, and valine.
- 6. The mutant of claim 2, wherein an amino acid residue at one or more of the positions 87, 98, 185, 244, 247, 295 or 297 is independently replaced by another amino acid residue.
- 7. The mutant of claim 3, wherein an amino acid residue at one or more of the positions 87, 98, 185, 244, 247, 295 or 297 is independently replaced by another amino acid residue.
- 8. A method of oxidizing a compound selected from the group consisting of a polycyclic aromatic hydrocarbon, a linear or branched alkane, a biphenyl compound or a halogenated variant thereof and a halogenated hydrocarbon, comprising the step of contacting said compound under oxidizing conditions with

mono-oxygenase cytochrome P-450.sub.cam wherein the tyrosine residue at position 96 is replaced by a small hydrophobic amino acid residue.

- 9. The method of claim 8, wherein the amino acid is selected from the group consisting of alanine, glycine, isoleucine, leucine, and valine.
- 10. The method of claim 8, wherein an amino acid residue at one or more of the positions 87, 98, 185, 244, 247, 295 or 297 is independently replaced by another amino acid residue.
- 11. The method of claim 9, wherein an amino acid residue at one or more of the positions 87, 98, 185, 244, 247, 295 or 297 is independently replaced by another amino acid residue.

WEST

Generate Collection

Print

Search Results - Record(s) 11 through 20 of 86 returned.

11. Document ID: US 6395299 B1

L11: Entry 11 of 86

File: USPT

May 28, 2002

US-PAT-NO: 6395299

DOCUMENT-IDENTIFIER: US 6395299 B1

TITLE: Matrices for drug delivery and methods for making and using the same

DATE-ISSUED: May 28, 2002

INVENTOR-INFORMATION:

NAME

CITY

STATE

ZIP CODE

COUNTRY

Babich; John W.

Scituate Syracuse MA NY

Zubieta; Jon Bonavia; Grant

Kensington

MD

US-CL-CURRENT: 424/484

Full Title Citation Front Review Classification Date Reference Sequences Attachments

Draw, Desc Image

KMC

12. Document ID: US 6388171 B1

L11: Entry 12 of 86

File: USPT

May 14, 2002

US-PAT-NO: 6388171

DOCUMENT-IDENTIFIER: US 6388171 B1

TITLE: Compositions and methods for fumonisin detoxification

DATE-ISSUED: May 14, 2002

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Duvick; Jon Des Moines IA
Maddox; Joyce Des Moines IA
Gilliam; Jacob Norwalk IA
Folkerts; Otto Guilford CT
Crasta; Oswald R. Branford CT

Full Title Citation Front Review Classification Date Reference Sequences Attachments

Draw Desc Image

KMAC

13. Document ID: US 6380145 B1

L11: Entry 13 of 86

File: USPT

Apr 30, 2002

US-PAT-NO: 6380145

DOCUMENT-IDENTIFIER: US 6380145 B1

TITLE: Cleaning compositions comprising a specific oxygenase

DATE-ISSUED: April 30, 2002

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Herbots; Ivan Maurice Alfons Jan B-1853 Strombeek-Bever BE

Barnabas; Mary Vijayarani Cincinnati OH 45061

Bettiol; Jean-Luc Philippe B-1853 Strombeek-Bever BE

Busch; Alfred B-1853 Strombeek-Bever BE

US-CL-CURRENT: 510/392; 510/114, 510/226, 510/300, 510/305, 510/306, 510/320, 510/374,

<u>510/530</u>



KWAC

14. Document ID: US 6365377 B1

L11: Entry 14 of 86

File: USPT

Apr 2, 2002

US-PAT-NO: 6365377

DOCUMENT-IDENTIFIER: US 6365377 B1

TITLE: Recombination of insertion modified nucleic acids

DATE-ISSUED: April 2, 2002

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Patten; Phillip A. Mountain View CA
Heinrichs; Volker Mountain View CA
Stemmer; Willem P. C. Los Gatos CA

US-CL-CURRENT: 435/91.1; 435/455, 435/463, 435/6, 436/94

Full: Title: Citation Front Review Classification Date Reference Sequences Attachments

Draw, Desc. Image

KWIC

15. Document ID: US 6365376 B1

L11: Entry 15 of 86

File: USPT

Apr 2, 2002

US-PAT-NO: 6365376

DOCUMENT-IDENTIFIER: US 6365376 B1

TITLE: Genes and enzymes for the production of adipic acid intermediates

DATE-ISSUED: April 2, 2002

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Brzostowicz; Patricia C. West Chester PA
Rouviere; Pierre E. Wilmington DE.

US-CL-CURRENT: 435/91.1; 435/252.3, 435/252.31, 435/252.32, 435/252.33, 435/252.35, 435/254.11, 435/254.2, 435/320.1, 435/91.2, 536/23.2, 536/23.7

Full : Title | Citation | Front | Review | Classification | Date | Reference | Sequences | Attachments | Draw, Desc | Image |

KWIC

16. Document ID: US 6310271 B1

L11: Entry 16 of 86 File: USPT Oct 30, 2001

US-PAT-NO: 6310271

DOCUMENT-IDENTIFIER: US 6310271 B1

TITLE: Polynucleotides encoding choline monooxygenase and plants transformed therewith

DATE-ISSUED: October 30, 2001

INVENTOR-INFORMATION:

NAME . CITY STATE ZIP CODE COUNTRY

Hanson; Andrew D. Gainesville FL Rathinasabapathi; Bala Gainesville FL

Burnet; Michael Les Hameaux FR

 $\begin{array}{l} \text{US-CL-CURRENT: } 800/278; \ 435/410, \ \underline{435/419}, \ 435/468, \ \underline{435/69.1}, \ \underline{536}/\underline{23.1}, \ \underline{536/23.6}, \\ \underline{800/285}, \ \underline{800/290}, \ \underline{800/295}, \ \underline{800}/\underline{306}, \ \underline{800}/\underline{312}, \ \underline{800/314}, \ \underline{800/317.2}, \ \underline{800}/\underline{317.3}, \ \underline{800}/\underline{317.3}, \ \underline{800}/\underline{317.4}, \\ \underline{800}/\underline{320.1}, \ \underline{800}/\underline{320.1}, \ \underline{800}/\underline{320.2}, \ \underline{800}/\underline{320.3}, \ \underline{800}/\underline{322} \\ \end{array}$

Full Title Citation Front Review Classification Date Reference Sequences Attachments

Draw Desc Image

KWIC

17. Document ID: US 6309883 B1

L11: Entry 17 of 86 File: USPT Oct 30, 2001

US-PAT-NO: 6309883

DOCUMENT-IDENTIFIER: US 6309883 B1

TITLE: Methods and compositions for cellular and metabolic engineering

DATE-ISSUED: October 30, 2001

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Minshull; Jeremy San Francisco CA Stemmer; Willem P. C. Los Gatos CA US-CL-CURRENT: 435/440; 435/6, 536/23.1, 536/24.3

Full Title Citation Front Review Classification Date Reference Sequences Attachments KWIC 18. Document ID: US 6300544 B1 L11: Entry 18 of 86 File: USPT Oct 9, 2001 US-PAT-NO: 6300544 DOCUMENT-IDENTIFIER: US 6300544 B1 TITLE: Cytochrome P450 monooxygenases DATE-ISSUED: October 9, 2001 INVENTOR-INFORMATION: CITY STATE ZIP CODE COUNTRY NAME DK Halkier; Barbara Ann Copenhagen K. Copenhagen N. DK Bak; Soren Kahn; Rachel Alice DK Copenhagen K. DK Moller; Birger Lindberg Bronshoj US-CL-CURRENT: 800/279; 435/183, 435/252.3, 435/320.1, 435/419, 435/6, 530/350, 530/370, 536/23.6, 536/24.1, 800/301Title Citation Front Review Classification Date Reference Sequences Attachments KAMC Draw, Desc | Image 19. Document ID: US 6268552 B1 L11: Entry 19 of 86 File: USPT Jul 31, 2001 US-PAT-NO: 6268552 DOCUMENT-IDENTIFIER: US 6268552 B1 TITLE: Transgenic seedless fruit comprising AGL or GH3 promoter operably linked to isopentenyl transferase or tryptophan monooxygenase coding DNA DATE-ISSUED: July 31, 2001 INVENTOR-INFORMATION: COUNTRY ZIP CODE NAME STATE CITY Li; Yi Mansfield Center US-CL-CURRENT: 800/317.4; 435/320.1, 800/278, 800/284, 800/298, 800/307, 800/308 KMC Full Title Citation Front Review Classification Date Reference Sequences Attachments Draw, Desc Image 20. Document ID: US 6255067 B1 Jul 3, 2001 L11: Entry 20 of 86 File: USPT

US-PAT-NO: 6255067

DOCUMENT-IDENTIFIER: US 6255067 B1

TITLE: cDNA encoding peptidyl-glycine alpha-amidating monooxygenase (PAM)

DATE-ISSUED: July 3, 2001

INVENTOR-INFORMATION:

NAME

CITY

ZIP CODE STATE

COUNTRY

DE

Keutmann; Henry T.

Concord Heidelberg

Schofield; Peter Rodriguez; Henry

Belmont

CA

MA

Eipper; Betty

Baltimore

MD

Mains; Richard

Baltimore

MD

 $\text{US-CL-CURRENT: } \underline{435}/\underline{69.1}; \ \underline{435}/\underline{252.3}, \ \underline{435}/\underline{254.2}, \ \underline{435}/\underline{320.1}, \ \underline{435}/\underline{325}, \ \underline{435}/\underline{349}, \ \underline{530}/\underline{350},$ <u>536/23.5</u>

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	KWIC
Draw	Desc	lmage								
		-								
	::::::::::::::::::::::::::::::::::::::				-	***********************		·····		
					Generate C			Print		
			***************************************	Ter					ocuments	
	<u> </u>				1115				ocuments	
monooxygenase.clm.									86	
				II.			: 4			

Display Format: CIT

Change Format

Previous Page

Next Page

WEST	
Generate Collection	Print

L15: Entry 2 of 20

File: USPT

Oct 29, 2002

US-PAT-NO: 6472191

DOCUMENT-IDENTIFIER: US 6472191 B1

TITLE: DNA FRAGMENT CARRYING TOLUENE MONOOXYGENASE GENE, RECOMBINANT PLASMID, TRANSFORMED MICROORGANISM, METHOD FOR DEGRADING CHLORINATED ALIPHATIC HYDROCARBON COMPOUNDS AND AROMATIC COMPOUNDS, AND METHOD FOR ENVIRONMENTAL REMEDIATION

DATE-ISSUED: October 29, 2002

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNȚRY Yano; Tetsuya Atsugi JP Nomoto; Tsuyoshi Komae JP Imamura; Takeshi Chigasaki JP

US-CL-CURRENT: 435/189; 435/252.3, 435/262.5, 435/320.1, 536/23.2

CLAIMS:

What is claimed is:

- 1. An isolated DNA fragment of about 5.3 Kb containing a toluene monoxygenase gene, having 3 BamHI, 1 ClaI, 1 EcoRI, 3 KpnI, 2 NcoI, 2 NspV, 2 ScaI, 2 SmaI, 2 SphI, 1 StuI, 0 DraI, 0 EcoRV, 0 HindIII, OHpaI, 0 NdeI, 0 PvuII, 0 ScaI, 0 Sse83871, 0 XbaI, 0 XhoI restriction sites, and having a restriction map of: ##STR3##
- 2. An isolated DNA fragment having a nucleotide sequence of SEQ ID NO: 1 in the Sequence Listing.
- 3. A DNA fragment encoding a protein having a toluene $\frac{\text{monooxygenase}}{\text{monooxygenase}}$ activity and being hybridizable to a nucleotide sequence from 200 . . . 4799 of SEQ ID NO: 1 or a complementary sequence thereof under stringent hybridization conditions.
- 4. A recombinant DNA comprising a vector which can replicate or can be maintained in a host and a DNA fragment according to any one of claims 1 to 3.
- 5. The recombinant DNA fragment according to claim 4, wherein the vector can be maintained or replicate in a bacterium.
- 6. A transformant obtainable by introducing into a host microorganism a recombinant DNA comprising the DNA fragment according to any one of claims 1 to 3 ligated to a vector which can replicate or be maintained in the host.
- 7. A method for producing a toluene monooxygenase, comprising: introducing into a host microorganism a recombinant DNA which comprises a vector which can replicate or can be maintained in the host microorganism and a DNA fragment according to any one of claims 1 to 3 to form a transformant which produces a toluene monooxygenase encoded by said DNA fragment.

- 8. A method for remedying an environment polluted with a pollutant being at least either of a halogenated aliphatic hydrocarbon compound or an aromatic compound, comprising a step of degrading the pollutant by bringing a transformant into contact with the pollutant, wherein the transformant is obtainable by introducing into a host microorganism a recombinant DNA constituted by ligating a vector which enables retention and replication in the host and a DNA fragment according to any one of claims 1 to 3.
- 9. The remediation method according to claim 8, wherein the environment is soil.
- 10. The remediation method according to claim 9 comprising the steps of: introducing an aqueous medium containing the transformant into the polluted soil; and supplying nutrients and/or oxygen for proliferation of the transformant in the polluted soil.
- 11. The remediation method according to claim 10 wherein the transformant is introduced in the soil by applying pressure through an injection well provided in the polluted soil.
- 12. The remediation method according to claim 9 wherein the polluted soil is introduced in a liquid phase containing the transformant.
- 13. The remediation method according to claim 9 wherein the polluted soil is brought into contact with a carrier holding the transformant.
- 14. The remediation method according to claim 8 wherein the environment is air.
- 15. The remediation method according to claim 14 wherein the polluted air is introduced into a liquid phase containing the transformant.
- 16. The remediation method according to claim 14 wherein the polluted air is brought into contact with a carrier holding the transformant.
- 17. The remediation method according to claim 16 wherein contact is carried out by placing the carrier holding the transformant in a container, introducing polluted air from one side of the container, and discharging cleaned air from another side.
- 18. The remediation method according to claim 8 wherein the halogenated aliphatic hydrocarbon compound is either trichloroethylene (TCE) or dichloroethylene (DCE).
- 19. The remediation method according to claim 8 wherein the aromatic compound is at least one of toluene, benzene, phenol, and cresol.

WEST	
Generate Collection	Print

L15: Entry 9 of 20 File: USPT Sep 28, 1999

US-PAT-NO: 5958757

DOCUMENT-IDENTIFIER: US 5958757 A

TITLE: Biological conversion of organic compounds

DATE-ISSUED: September 28, 1999

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Steffan; Robert Jon Newtown PA
McClay; Kevin Rock Morrisville PA

US-CL-CURRENT: 435/262.5; 435/170, 435/264, 435/289.1, 435/874, 570/220, 588/248

CLAIMS:

We claim:

- 1. A method for oxidizing saturated aliphatic halocarbons comprising contacting said halocarbons with an aromatic oxygenase capable of oxidizing said halocarbons, which is produced by a microorganism wherein said saturated halocarbon is selected from the group consisting of chloroform, bromoform, 1,2-dichloroethane, 1,2 dibromoethane, monochloroethane, and monobromoethane.
- 2. The method of claim 1 wherein said saturated aliphatic halocarbon is chloroform.
- 3. The method of claim 1 which further comprises providing a co-substrate to support degradation of said halocarbon by said microorganism.
- 4. The method of claim 3 wherein said co-substrate is selected from the group consisting of toluene, phenol, benzene, ethylbenzene, and xylene.
- 5. The method of claim 1 wherein said saturated halocarbon is contacted with said microorganism in water.
- 6. The method of claim 1 wherein said saturated halocarbon is contacted with said microorganism in soil.
- 7. The method of claim 1 wherein said saturated halocarbon is contacted with said microorganism in vapor phase.
- 8. The method of claim 1 wherein said saturated halocarbon is contacted with said aromatic oxygenase-producing microorganism within a bioreactor.
- 9. The method of claim 8 wherein said bioreactor is a fixed film bioreactor.
- 10. The method of claim 8 wherein said bioreactor is a suspended growth bioreactor.
- 11. The method of claim 1 wherein said saturated halocarbon is contacted with

said aromatic oxygenase-producing bacteria in situ.

- 12. The method of claim 11 wherein said saturated halocarbon is present in soil or sludge.
- 13. The method of claim 11 wherein said saturated halocarbon is present in groundwater.
- 14. The method of claim 1 wherein said aromatic oxygenase is a toluene monooxygenase.
- 15. The method of claim 1 wherein said aromatic oxygenase-producing microorganism is selected from the group consisting of Pseudomonas mendocina KR1, ATCC 55706; Strain ENVPC5; and Strain ENVBF1 ATCC 55819.
- 16. The method of claim 1 wherein said aromatic oxygenase-producing microorganism is a recombinant microorganism consisting of a host microorganism containing cloned aromatic oxygenase genes.
- 17. A method for oxidizing saturated aliphatic halocarbons comprising contacting said halocarbons with an aromatic oxygenase capable of oxidizing said halocarbons, which is produced by an aerobic bacteria wherein said saturated halocarbon is selected from the group consisting of chloroform, bromoform, 1,2-dichloroethane, 1,2 dibromoethane, monochloroethane, and monobromoethane.
- 18. A method for oxidizing saturated aliphatic halocarbons comprising contacting said halocarbons with a toluene-4-monooxygenase which is produced by an aerobic bacteria.

WEST Search History

DATE: Tuesday, January 14, 2003

Set Name side by side	Query	Hit Count	Set Name result set
DB=US	PT; PLUR=YES; OP=ADJ		
L17	L13 and aromatic sibstrate	0	L17
L16	L13 and halo aromatic sibstrate	0	L16
L15	111 and oxidation and benzene	20	L15
L14	111 and oxidation and dichlorobenzene	0	L14
L13	111 and oxidation	51	L13
L12	mutant monooxygenase.clm.	0	L12
L11	monooxygenase.clm.	86	L11
L10	mono-oxygenase.clm.	9	L10
L9	mutant mono-oxygenase	2	L9
L8	oxidizing dichlorophenyl	0	L8
L7	oxidizing pentachlorophenyl	0	L7
L6	L1 and pentachlorophenyl	1	L6
L5	L1 and dichlorophenyl	7	L5
L4	L2 and pentachlorobiphenyl	0	L4
L3	L2 and dichlorobiphenyl	0	L3
L2	L1 and dichlorobenzene	20	L2
L1	monooxygenase	653	L1

END OF SEARCH HISTORY

WEST Search History

DATE: Tuesday, January 14, 2003

Set Name side by side	Query	Hit Count	Set Name result set
DB=US	SPT; PLUR=YES; OP=ADJ		
L21	L20 and p-450	3	L21
L20	oxidizing and monooxygenase.clm.	33	L20
L19	oxidizing and monooxygenase	193	L19
L18	method for oxidizing and monooxygenase	0	L18
L17	L13 and aromatic sibstrate	0	L17
L16	L13 and halo aromatic sibstrate	0	L16
L15	111 and oxidation and benzene	20	L15
L14	111 and oxidation and dichlorobenzene	0	L14
L13	111 and oxidation	51	L13
L12	mutant monooxygenase.clm.	0	L12
L11	monooxygenase.clm.	86	L11
L10	mono-oxygenase.clm.	9	L10
L9	mutant mono-oxygenase	2	L9
L8	oxidizing dichlorophenyl	0	L8
L7	oxidizing pentachlorophenyl	0	L7
L6	L1 and pentachlorophenyl	1	L6
L5	L1 and dichlorophenyl	7	L5
L4	L2 and pentachlorobiphenyl	0	L4
L3	L2 and dichlorobiphenyl	0	L3
L2	L1 and dichlorobenzene	20	L2
L1	monooxygenase	653	L1

END OF SEARCH HISTORY

1/14/03 1:49 PM

WEST

End of Result Set

Generate Collection Print

L5: Entry 1 of 1

File: DWPI

May 7, 1997

DERWENT-ACC-NO: 1997-229326

DERWENT-WEEK: 200046

COPYRIGHT 2003 DERWENT INFORMATION LTD

TITLE: Mutant of the mono-oxygenase cytochrome P-450cam - useful in catalysis of oxidation of a wide range of organic substrates, such as lindane.

INVENTOR: FLITSCH, S L; HART, A J; NICKERSON, D P; WONG, L

PRIORITY-DATA: 1995WO-GB02588 (November 2, 1995), 1995GB-0022407 (November 1, 1995)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
GB 2306485 A	May 7, 1997		040	C12N009/02
US 6117661 A	September 12, 2000		000	C12N009/02
WO 9716553 A1	May 9, 1997		000	C12N015/53
AU 9673236 A	May 22, 1997		000	C12N015/53
GB 2306485 B	December 9, 1998		000	C12N009/02
CZ 9801273 A3	January 13, 1999		000	C12N015/53
EP 906431 A1	April 7, 1999	E	000	C12N015/53
SK 9800555 A3	April 13, 1999		000	C12N015/53
CN 1212015 A	March 24, 1999		000	C12N015/53
NZ 320497 A	September 29, 1999		000	C12N015/53
AU 716583 B	March 2, 2000		000	C12N015/53
JP 2000508163 W	July 4, 2000		039	C12N009/02

INT-CL (IPC): C12 N 9/02; C12 N 15/00; C12 N 15/09; C12 N 15/53; C12 N 15/78; C12 P 7/02; C12 P 7/22; C12 R 1:19; C12 R 1:19

ABSTRACTED-PUB-NO: GB 2306485A

BASIC-ABSTRACT:

Mutant of the mono-oxygenase cytochrome P-450cam, in which the cysteine residue at position 334 is removed, is new.

USE- Mono-oxygenase cytochrome P-450cam from P. putida catalyses the regio- and stereoselective hydroxylation of camphor to 5-exo-hydroxycamphor. The mutant enzyme may be used to catalyse the oxidation of a relatively wide range of organic substrates (such as lindane), whether or not these are substrates for the wild-type protein.

ADVANTAGE- A C334A mutant of P-450cam did not show any evidence of aggregation even at mM concentrations at room temperature over a period of three days. This property is expected to improve protein handling, storage and increased catalyst lifetime.

ABSTRACTED-PUB-NO:

GB 2306485B

EQUIVALENT-ABSTRACTS:

Mutant of the mono-oxygenase cytochrome P-450cam, in which the cysteine residue at position 334 is removed, is new.

USE- Mono-oxygenase cytochrome P-450cam from P. putida catalyses the regio- and stereoselective hydroxylation of camphor to 5-exo-hydroxycamphor. The mutant enzyme may be used to catalyse the oxidation of a relatively wide range of organic substrates (such as lindane), whether or not these are substrates for the wild-type protein.

ADVANTAGE- A C334A mutant of P-450cam did not show any evidence of aggregation even at mM concentrations at room temperature over a period of three days. This property is expected to improve protein handling, storage and increased catalyst lifetime.

US 6117661A

Mutant of the mono-oxygenase cytochrome P-450cam, in which the cysteine residue at position 334 is removed, is new.

USE- Mono-oxygenase cytochrome P-450cam from P. putida catalyses the regio- and stereoselective hydroxylation of camphor to 5-exo-hydroxycamphor. The mutant enzyme may be used to catalyse the oxidation of a relatively wide range of organic substrates (such as lindane), whether or not these are substrates for the wild-type protein.

ADVANTAGE- A C334A mutant of P-450cam did not show any evidence of aggregation even at mM concentrations at room temperature over a period of three days. This property is expected to improve protein handling, storage and increased catalyst lifetime.

ABSTRACTED-PUB-NO: GB 2306485A EQUIVALENT-ABSTRACTS: GB 2306485B Mutant of the mono-oxygenase cytochrome P-450cam, in which the cysteine residue at position 334 is removed, is new. USE- Mono-oxygenase cytochrome P-450cam from P. putida catalyses the regio- and stereoselective hydroxylation of camphor to 5-exo-hydroxycamphor. The mutant enzyme may be used to catalyse the oxidation of a relatively wide range of organic substrates (such as lindane), whether or not these are substrates for the wild-type protein. ADVANTAGE- A C334A mutant of P-450cam did not show any evidence of aggregation even at mM concentrations at room temperature over a period of three days. This property is expected to improve protein handling, storage and increased catalyst lifetime. US 6117661A Mutant of the mono-oxygenase cytochrome P-450cam, in which the cysteine residue at position 334 is removed, is new. USE- Mono-oxygenase cytochrome P-450cam from P. putida catalyses the regio- and stereoselective hydroxylation of camphor to 5-exo-hydroxycamphor. The mutant enzyme may be used to catalyse the oxidation of a relatively wide range of organic substrates (such as lindane), whether or not these are substrates for the wild-type protein. ADVANTAGE- A C334A mutant of P-450cam did not show any evidence of aggregation even at mM concentrations at room temperature over a period of three days. This property is expected to improve protein handling, storage and increased catalyst lifetime.

CHOSEN-DRAWING: Dwg.0/2



End of Result Set

Generate Collection Print

L7: Entry 2 of 2

File: DWPI

May 8, 1996

DERWENT-ACC-NO: 1996-211678

DERWENT-WEEK: 200112

COPYRIGHT 2003 DERWENT INFORMATION LTD

TITLE: New mutant forms of monooxygenase cytochrome P-450 (cam) - with amino acid substitutions at positions 96 and 334, useful for oxidn. of wide range of aliphatic and aromatic, opt. halogenated, hydrocarbon(s)

INVENTOR: FLITSCH, S L; NICKERSON, D P; WONG, L; WONG, L L; NICKERSON, D

PRIORITY-DATA: 1994GB-0022205 (November 3, 1994)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
GB 2294692 A	May 8, 1996		042	C12N009/02
KR 234348 B1	December 15, 1999		000	C12N015/53
WO 9614419 A1	May 17, 1996	E	044	C12N015/53
AU 9538117 A	May 31, 1996		000	C12N015/53
EP 789770 A1	August 20, 1997	E	000	C12N015/53
CZ 9701277 A3	October 15, 1997		000	C12N015/53
SK 9700545 A3	February 4, 1998		000	C12N015/53
JP 10503658 W	April 7, 1998		039	C12N015/09
NZ 294904 A	September 24, 1998		000	C12P007/02
KR 97707288 A	December 1, 1997		000	C12N015/53
GB 2294692 B	January 20, 1999		000	C12N009/02
AU 705736 B	June 3, 1999		000	C12N015/53
RU 2133774 C1	July 27, 1999		000	C12N015/53
US 6100074 A	August 8, 2000		000	C12N009/02

ABSTRACTED-PUB-NO: GB 2294692A

BASIC-ABSTRACT:

New mutants (I) of mono-oxygenase cytochrome P-450 can have 96 Tyr and/or 334 Cys replaced by any other amino acid (aa) provided: (a) replacement is not by Phe or (b) the mutant is able to oxidise any of polycyclic aromatic hydrocarbons, branched or linear alkanes, diphenyls or biphenyls (including halogenated derivs.) or halogenated hydrocarbons.

USE - (I) are used to oxidise the substrates named above, e.g. to produce starting materials (partic. in homo-chiral form) for organic synthesis.

ADVANTAGE - (I) can oxidise a wide range of substrates (claimed) (alteration of the aromatic binding site pocket reduces specificity for camphor, the natural substrate). Elimination of 334 Cys improves stability and prevents unwanted dimerisation during purificn.

ABSTRACTED-PUB-NO:

GB 2294692B EQUIVALENT-ABSTRACTS:

New mutants (I) of mono-oxygenase cytochrome P-450 can have 96 Tyr and/or 334 Cys replaced by any other amino acid (aa) provided: (a) replacement is not by Phe or (b) the mutant is able to oxidise any of polycyclic aromatic hydrocarbons, branched or linear alkanes, diphenyls or biphenyls (including halogenated derivs.) or halogenated hydrocarbons.

USE - (I) are used to oxidise the substrates named above, e.g. to produce starting materials (partic. in homo-chiral form) for organic synthesis.

ADVANTAGE - (I) can oxidise a wide range of substrates (claimed) (alteration of the aromatic binding site pocket reduces specificity for camphor, the natural substrate). Elimination of 334 Cys improves stability and prevents unwanted dimerisation during purificn.

US 6100074A

New mutants (I) of mono-oxygenase cytochrome P-450 can have 96 Tyr and/or 334 Cys replaced by any other amino acid (aa) provided: (a) replacement is not by Phe or (b) the mutant is able to oxidise any of polycyclic aromatic hydrocarbons, branched or linear alkanes, diphenyls or biphenyls (including halogenated derivs.) or halogenated hydrocarbons.

USE - (I) are used to oxidise the substrates named above, e.g. to produce starting materials (partic. in homo-chiral form) for organic synthesis.

ADVANTAGE - (I) can oxidise a wide range of substrates (claimed) (alteration of the aromatic binding site pocket reduces specificity for camphor, the natural substrate). Elimination of 334 Cys improves stability and prevents unwanted dimerisation during purificn.

ABSTRACTED-PUB-NO: GB 2294692A EQUIVALENT-ABSTRACTS: GB 2294692B New mutants (I) of mono-oxygenase cytochrome P-450 can have 96 Tyr and/or 334 Cys replaced by any other amino acid (aa) provided: (a) replacement is not by Phe or (b) the mutant is able to oxidise any of polycyclic aromatic hydrocarbons, branched or linear alkanes, diphenyls or biphenyls (including halogenated derivs.) or halogenated hydrocarbons. USE - (I) are used to oxidise the substrates named above, e.g. to produce starting materials (partic. in homo-chiral form) for organic synthesis. ADVANTAGE - (I) can oxidise a wide range of substrates (claimed) (alteration of the aromatic binding site pocket reduces specificity for camphor, the natural substrate). Elimination of 334 Cys improves stability and prevents unwanted dimerisation during purificn. US 6100074A New mutants (I) of mono-oxygenase cytochrome P-450 can have 96 Tyr and/or 334 Cys replaced by any other amino acid (aa) provided: (a) replacement is not by Phe or (b) the mutant is able to oxidise any of polycyclic aromatic hydrocarbons, branched or linear alkanes, diphenyls or biphenyls (including halogenated derivs.) or halogenated hydrocarbons. USE - (I) are used to oxidise the substrates named above, e.g. to produce starting materials (partic. in homo-chiral form) for organic synthesis. ADVANTAGE - (I) can oxidise a wide range of substrates (claimed) (alteration of the aromatic binding site pocket reduces specificity for camphor, the natural substrate). Elimination of 334 Cys improves stability and prevents unwanted dimerisation during purificn.

CHOSEN-DRAWING: Dwg.0/2

WEST

Generate Collection

Print

Search Results - Record(s) 1 through 2 of 2 returned.

1. Document ID: US 6117661 A

L15: Entry 1 of 2

File: USPT

Sep 12, 2000

US-PAT-NO: 6117661

DOCUMENT-IDENTIFIER: US 6117661 A

TITLE: Mutant mono-oxygenase cytochrome P450cam

DATE-ISSUED: September 12, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Wong; Luet-Lok Oxford GB
Flitsch; Sabine Lahja Edinburgh GB
Nickerson; Darren Paul Oxford GB
Hart; Alwyn James Loughborough GB

US-CL-CURRENT: 435/189; 435/252.3, 435/320.1, 435/471, 435/69.1, 536/23.2

Full | Title | Citation | Front | Review | Classification | Date | Reference | Sequences | Attachments | Claims | KMC | Draw, Desc | Image |

2. Document ID: US 6100074 A

L15: Entry 2 of 2

File: USPT

Aug 8, 2000

US-PAT-NO: 6100074

DOCUMENT-IDENTIFIER: US 6100074 A

TITLE: Mutant mono-oxygenase cytochrome P-450 .sub.cam

DATE-ISSUED: August 8, 2000

INVENTOR-INFORMATION:

NAME CITY STATE ZIP CODE COUNTRY

Flitsch; Sabine Lahja Edinburgh GB
Nickerson; Darren Paul Oxford GB
Wong; Luet-Lok Oxford . GB

US-CL-CURRENT: 435/189; 435/132, 435/69.1, 530/402

Full Title Citation Front Review Classification Date Reference Sequences Attachments Claims KMC
Draw Desc Image

WEST Search History

DATE: Tuesday, January 14, 2003

Hit Count	Set Name result set
2	L15
0	L14
. 0	L13
0	L12
0	L11
0	L10
. 0	L9
ome 0	L8
2	L7
1	L6
1	L.5
0	L4
20	L3
1	L2
0	L1
	0 0 0 0 0 0 0 0 0 0 2 1

END OF SEARCH HISTORY